arXiv:1205.1549v1 [hep-th] 7 May 2012

Vacuum states on timelike hypersurfaces in quantum field theory[*]

Ralf Banisch[†]
Freie Universität Berlin

Frank Hellmann[‡]
Albert Einstein Institute

Dennis Rätzel[§]
Albert Einstein Institute

March 3, 2013

We show how to adapt the canonical arguments for choosing the physically correct vacuum state to the case of quantum field theory with states on timelike hypersurfaces as formulated in the general boundary formalism. We demonstrate that these fail for evanescent modes which occur naturally in quantum field theory in the presence of the equivalent of a dielectric boundary. We explicitly study the response of an Unruh-DeWitt type detector to these modes. Using this we find that the physically correct vacuum state has to depend on the physical situation outside of the boundaries of the spacetime region considered. Thus it cannot be determined by general principles pertaining only to the spacetime region under consideration.

PACS numbers: 11.10.-z, 04.62.+v

1. Introduction

The general boundary formalism for quantum field theory is a formalism that allows to consider quantum field theories on arbitrary hypersurfaces in spacetime. This formalism was first proposed axiomatically in [Oec05, Oec06, Oec07], in analogy to the axiomatic formulation of topological quantum field theories due to Atiyah [Ati89].

In recent years this formalism has seen a wealth of technical development. Quantization techniques for ordinary quantum field theory have been adapted to this formalism and rigorous constructions for linear and affine field theories have been given [CO08a, CO08b, Oec08, CO09, Oec10, CDO11, CO11, Oec12b, Oec12a, Oec11, Oec12c]. The formalism was used to investigate the behavior of the S-matrix at spatial infinity [CO08a, CO08b, CDO11] and the general notion of observables for these theories was clarified in [Oec12b].

In this paper we study in detail the physical and symmetry conditions one can impose to determine the remaining ambiguities in the quantization, that is, the choice of vacuum state, or in the language of holomorphic quantization, the complex structure.

---

[*] AEI-2012-040

[†] ralf.banisch@fu-berlin.de

[‡] frank.hellmann@aei.mpg.de

[§] dennis.raetzel@aei.mpg.de

We consider two types of conditions. The first is based on adapting the Ashtekar-Magnon condition [AM75] to timelike hypersurfaces. The second is based on coupling an Unruh-DeWitt detector moving along a specified world line and studying the transition rates given certain boundary states.

We find that for the ordinary field modes, the propagating waves, the Ashtekar-Magnon like condition is satisfactory and produces the results of the canonical case. Furthermore we show that Ashtekar-Magnon like conditions can actually subsume the symmetry considerations ordinarily used. Using the detector model we can also show that the condition that a detector at rest does not click in vacuum can also reproduce the complex structure singled out by the Ashtekar-Magnon condition.

For field modes that behave exponential in space, the evanescent waves, an Ashtekar-Magnon like condition cannot be satisfied under the given assumptions. In the case of the detector model it is also less clear how to select the boundary states for the no-click condition. Thus we also study the actual transition rates in a concrete experimental set up capable of producing modes that are exponentially decaying in space [CM71, CMD72].

This is done by putting a dielectric boundary outside of the timelike region we consider. We find that we can determine a unique unitary complex structure by requiring the transition rates for various boundary states to reproduce the experimentally correct result, and that the complex structure depends on the details of the physics outside the spacetime region under consideration. We also find that non-unitary complex structures are necessary to satisfy Ashtekar-Magnon like conditions asymptotically, where they can be applied again.

1.1. An outline of the paper

In order to make the paper largely self contained we begin by giving an overview of the pertinent structures of a general boundary quantum field theory in section 2. We focus less on the mathematically rigorous constructions and more on clarifying the structural ingredients for the interpretation of the theory.

In section 3 we review in detail how to construct a general boundary quantum field theory with timelike hypersurfaces. We begin by briefly discussing the Schrödinger-Feynman quantization in section 3.1. We then introduce the language of holomorphic quantization in 3.2, discussing in particular the notion of unitarity in the theory, and the construction of the one particle and coherent states. We pay particular attention to the quantization ambiguity given by the choice of complex structure.

In section 4 we discuss the Ashtekar-Magnon condition [AM75] for choosing complex structures for canonical quantization on arbitrary spacetimes. After reviewing the condition for spacelike hypersurfaces in section 4.1, we demonstrate in 4.2 that this condition can also be adapted to reproduce symmetry principles for arbitrary hyperplanes in a straightforward way. We then discuss the case of timelike hypersurfaces in 4.3. For the sector of the theory dealing with propagating waves the equivalent of the Ashtekar-Magnon condition discussed in 4.3.1 does determine a natural complex structure. We show that this condition cannot be applied to the case of evanescent waves. In 4.3.2 we discuss in detail how to interpret the results of the preceding sections physically and how the physical role of choosing a boundary state and the complex structure change vis-a-vis the canonical theory.

In section 5 we study in detail how to couple a detector in the language of the general boundary formalism and how to use the dependence of the detection rate on the complex structure to determine the latter in physical terms. We begin by developing perturbation theory in the holomorphic language in section 5.1. Using this perturbation theory we show how to couple an Unruh-DeWitt type detector to a quantum field theory in the general boundary formulation in section 5.2, which to our knowledge has not been studied in the literature so far. We show that, using symmetry arguments and an appropriate choice of boundary states, we can determine the complex structure for propagating modes by requiring a detector at rest to not click in section 5.3, thus bypassing the Ashtekar-Magnon condition. As the appropriate choice of boundary state is unclear for evanescent waves we then consider an actual physical example in which evanescent waves appear and describe it in the general boundary formulation in section 5.4. This example is a toy model of a free electromagnetic field in the presence of a dielectric boundary. The excitation rates induced by evanescent waves in this case have been studied theoretically and experimentally in [CM71, CMD72]. We show that in order to reproduce the transition rates while maintaining the ability to model the two situations of particle emission and particle absorption as orthogonal subsets in the boundary Hilbert space, which is the situation one has in canonical quantization, the complex structure has to depend on the physical situation at the dielectric boundary, even if this boundary is outside of the spacetime region considered. We conclude that a complex structure for evanescent waves cannot be chosen by general physical arguments pertaining only to the spacetime region under consideration. We see that a unitary complex structure exists satisfying our requirements, but that it doesn't satisfy Ashtekar-Magnon like conditions asymptotically, where they apply. These can be satisfied by a choice of non-unitary complex structure. Such a non-unitary complex structure is used for the first time in interacting GBQFT in this article.

2. The GB Formulation of quantum field theories

The general boundary formulation of quantum field theories (henceforth called GBF or GBQFT) is a framework that was introduced in [Oec08, Oec10]. In [Oec12c], the construction with the complete set of axioms spelled out in detail can be found in a top-down approach: The axioms of the GBF are given, and then certain theories satisfying these axioms are constructed. In the following we shall only give an introduction to the basic ideas and constructions of GBQFT in a bottom-up approach somewhat complementary to the discussions given in the references.

In the usual framework of quantum theory, one is interested in the transition amplitude $\rho(\psi_{t_1}, \psi_{t_2})$ that a certain state ψ_{t_1} in the Hilbert space $\mathcal{H}_{t_1}$ at time t_1 evolves into some other state ψ_{t_2} in the Hilbert space at time t_2. In other words,

$$\rho : \mathcal{H}_{t_1} \otimes \mathcal{H}_{t_2}^* \to \mathbb{C},$$

where $\mathcal{H}_{t_2}^*$ is the dual Hilbert space. These initial and final Hilbert spaces are assigned to the instants of time t_1 and t_2. In QFT these instants of time become equal-time or Cauchy hypersurfaces Σ_1 and Σ_2, and one is again interested in the transition amplitudes between states on pairs of Cauchy hypersurfaces. Obviously there is a certain region M (subset of spacetime) with boundary $\partial M = \Sigma_1 \cup \bar{\Sigma}_2$ (the bar denotes change of orientation). We may construct the boundary Hilbert space $\mathcal{H}_{\partial M} = \mathcal{H}_{t_1} \otimes \mathcal{H}_{t_2}^*$ and view ρ as a map from $\mathcal{H}_{\partial M}$ to $\mathbb{C}$. The idea of GBQFT is to generalize this to arbitrary spacetime regions M with arbitrary boundaries ∂M. To this end, let us fix some language. d is the dimension of spacetime. We make the following definitions:

- A *region* is an oriented d-dimensional topological manifold, possibly with boundary.
- A *hypersurface* is an oriented $(d-1)$-dimensional topological manifold without boundary.
- If Σ is a hypersurface, then $\bar{\Sigma}$ denotes the same manifold with opposite orientation.

Such a collection of regions and hypersurfaces is called a *spacetime system.* A GBQFT on a given spacetime system is a model satisfying a number of axioms similar to those of a topological quantum field theory [Ati89]. We outline the most important ones now, omitting all technical conditions needed to make the theory well defined. They can be found in full detail in [Oec12c].

1. Associated with each hypersurface Σ is a complex separable Hilbert space $\mathcal{H}_\Sigma$ with inner product $\langle \cdot, \cdot \rangle_\Sigma$ called the *state space* of Σ. There is a conjugate linear isometry $\iota_\Sigma : \mathcal{H}_\Sigma \to \mathcal{H}_{\bar{\Sigma}}$. This map is an involution in the sense that $\iota_{\bar{\Sigma}} \circ \iota_\Sigma$ is the identity on $\mathcal{H}_\Sigma$.
2. If Σ decomposes into a disjoint union of hypersurfaces, $\Sigma = \Sigma_1 \cup \ldots \cup \Sigma_n$, then $\mathcal{H}_\Sigma$ is isometrically isomorphic to $\mathcal{H}_{\Sigma_1} \otimes \ldots \otimes \mathcal{H}_{\Sigma_n}$. We shall always write $\mathcal{H}_\Sigma = \mathcal{H}_{\Sigma_1} \otimes \ldots \otimes \mathcal{H}_{\Sigma_n}$.

3. Associated with each region M there is a linear map $\rho_M : \mathcal{H}^\circ_{\partial M} \to \mathbb{C}$ where $\mathcal{H}^\circ_{\partial M}$ is dense in $\mathcal{H}_{\partial M}$. This is called the *amplitude map*. It has to satisfy a number of gluing properties.

To construct a GBQFT means to construct the state spaces $\mathcal{H}_\Sigma$ and the amplitude maps ρ_M. Our preferred way of doing this is by quantization of a classical field theory. Standard quantization techniques usually require spacetime to be foliated into Cauchy hypersurfaces, so they have to be generalized. In the following, we shall review the classical field theory setting in a geometric fashion which is more apt for constructing a GBQFT. For technical simplicity, only the case of linear field theory, for which the solution spaces are linear spaces is considered. We will mostly consider free fields which fall into this category. In the next section, we shall review two quantization settings discussed in the literature [CO08a, CO08b, Oec08, CO09, Oec10, CDO11, CO11, Oec12b, Oec12a, Oec11, Oec12c].

The classical scalar field theory starts with an action leading to first order Euler-Lagrange equations. For any region M, define

$$S_M(\phi) := \int_M dV L(\phi(x), \partial\phi(x))$$

where $L(\phi, \partial\phi, x)$ is the usual Lagrangian density for the Klein-Gordon field on a Lorentzian spacetime

$$L(\phi, \partial\phi, x) := \frac{1}{2} g^{\mu\nu} \partial_\mu \phi(x) \partial_\nu \phi(x) - \frac{1}{2} m^2 \phi^2(x)\,, \tag{1}$$

and $dV = \sqrt{-\det g}\, \epsilon_{\mu_1 \cdots \mu_4} dx^{\mu_1} \cdots dx^{\mu_4}$ is the volume form on M. Denote by L_Σ the real vector space of germs [1] of solutions to the Euler-Lagrange-equations in a neighborhood of Σ. If Σ is a Cauchy surface, then obviously every element of L_Σ can be uniquely extended to a solution everywhere in the region. But even if this is not the case, L_Σ contains enough information to extend analytic functions uniquely. In this paper we will always deal with analytic functions.

From the classical action derives the symplectic potential

$$(\theta_\Sigma)_\phi(X) := \int_\Sigma d\sigma_\mu X(x) \frac{\partial L}{\partial(\partial_\mu \phi)}(x)$$

where $\phi \in L_\Sigma$ is a solution, and X is a tangent vector to ϕ, i.e. $X \in T_\phi L_\Sigma \approx L_\Sigma$ [2] and $d\sigma_\mu = \sqrt{-\det g}\, \epsilon_{\mu_1 \cdots \mu_3 \mu} dx^{\mu_1} \cdots dx^{\mu_3}$. From this we then immediately have the bilinear map

$$[\cdot, \cdot]_\Sigma : L_\Sigma \times L_\Sigma \to \mathbb{R}, \quad [\phi, \xi]_\Sigma := (\theta_\Sigma)_\phi(\xi)\,,$$

and we obtain an antisymmetric bilinear map by anti-symmetrization:

$$\omega_\Sigma : L_\Sigma \times L_\Sigma \to \mathbb{R}, \quad \omega_\Sigma(\phi, \phi') = \frac{1}{2}[\phi, \phi']_\Sigma - \frac{1}{2}[\phi', \phi]_\Sigma$$

which we call the symplectic form.

If Σ decomposes into a disjoint union of hypersurfaces, $\Sigma = \Sigma_1 \cup \ldots \cup \Sigma_n$, then we obviously have $L_\Sigma = L_{\Sigma_1} \oplus \ldots \oplus L_{\Sigma_n}$, and $\theta_\Sigma = \theta_{\Sigma_1} + \ldots + \theta_{\Sigma_n}$.

For a region M we define $\omega_{\partial M}$ as the symplectic form associated to the boundary ∂M. We find that for two solutions ϕ, ϕ' of the Euler-Lagrange equations to (1) the symplectic form vanishes. This is because, using the specific form of (1), we have

$$\begin{aligned} \omega_{\partial M}(\phi, \phi') &= \int_{\partial M} d\sigma_\mu \left(\phi(x) \frac{\partial L}{\partial(\partial_\mu \phi')}(x) - \phi'(x) \frac{\partial L}{\partial(\partial_\mu \phi)}(x) \right) \\ &= \int_M dV \partial_\mu \left(\phi(x) \frac{\partial L}{\partial(\partial_\mu \phi')}(x) - \phi'(x) \frac{\partial L}{\partial(\partial_\mu \phi)}(x) \right) = 0\,. \end{aligned} \tag{2}$$

---

[1] For functions f and g on a topological space X, define $f \sim_x g$ if there is a neighborhood U of x such that $f|_U = g|_U$. The equivalence classes of $\sim_x$ are called *germs at x*. In particular, $f \sim_x g$ implies that all derivatives of f and g at x are equal.

[2] $T_\phi L_\Sigma$ is L_Σ linearized around ϕ, but since L_Σ is assumed to be linear, we can set $T_\phi L_\Sigma \simeq L_\Sigma$

3. On Quantizations and Complex Structures

We will consider two different quantization schemes discussed in the literature here. We first briefly introduce Schrödinger-Feynman quantization [Oec11] in the language given above and then focus mostly on holomorphic quantization, which is a special case of geometric quantization. In both cases, an additional datum is needed. It is well known [Woo97] that the additional datum needed for holomorphic quantization is a choice of a compatible and positive complex structure, i.e. a linear map $J_\Sigma : L_\Sigma \to L_\Sigma$ such that $J_\Sigma^2 = -\mathrm{id}_\Sigma$, $\omega_\Sigma(J_\Sigma\cdot, J_\Sigma\cdot) = \omega_\Sigma(\cdot,\cdot)$ and $\omega_\Sigma(\cdot, J_\Sigma\cdot)$ positive definite. That turns L_Σ into a complex Hilbert space and allows the definition coherent states that form the dense subset required in axiom 3. The additional datum needed for Schrödinger-Feynman-quantization is a vacuum state. It will turn out that there is a one-to-one correspondence between the choice of a complex structure and that of a vacuum state [Oec12c].

3.1. Schrödinger-Feynman Quantization

The basic idea here is to construct the Hilbert space as wave functions on the *configuration space* associated to the hypersurface Σ. This is constructed as follows: Define the subspaces $M_\Sigma, N_\Sigma \in L_\Sigma$ as

$$\begin{aligned} M_\Sigma &:= \{\tau \in L_\Sigma : [\xi,\tau]_\Sigma = 0 \forall \xi \in L_\Sigma\} \\ N_\Sigma &:= \{\tau \in L_\Sigma : [\tau,\xi]_\Sigma = 0 \forall \xi \in L_\Sigma\} \end{aligned}$$

We can think of M_Σ as the space of momenta (those solutions where field values vanish on Σ while their derivatives do not), and N_Σ as the space of 'positions' (vice versa). The configuration space Q_Σ is then the quotient space

$$Q_\Sigma := L_\Sigma / M_\Sigma$$

with quotient map $q_\Sigma : L_\Sigma \to Q_\Sigma$. To determine the Hilbert space $\mathcal{H}_\Sigma^S$ of the Schrödinger representation one needs the additional datum of a vacuum state [Oec12c]. This can be encoded [Oec11] into a symmetric bilinear form $\Omega_\Sigma : Q_\Sigma \times Q_\Sigma \to \mathbb{C}$ with positive definite real part. The vacuum state is then given by the wave function $K_0^S : Q_\Sigma \to \mathbb{C}$ given by

$$K_0^S(\phi) = \exp\left(-\frac{1}{2}\Omega(\phi,\phi)\right). \tag{3}$$

3.2. Holomorphic Quantization

We now turn to holomorphic quantization. For the current context this quantization procedure was given in detail in [Oec10],[Oec12a] for linear and affine field theories. It is a special case of geometric quantization [Woo97]. We outline it here, again ignoring all technical difficulties.

The basic idea of geometric quantization is to rigorously define the textbook canonical quantization procedure in cases where the configuration spaces of interest are smooth symplectic manifolds, as explained above. Textbook quantization tells you to proceed by expanding the classical solutions into Fourier modes and then dividing them up into two sets: Positive and negative frequency modes. The coefficients in front of the positive and negative frequency modes are then turned into creation and annihilation operators. The notion of positive and negative frequency is meant relative to a nowhere vanishing timelike, hypersurface orthogonal Killing vector field $\frac{\partial}{\partial t}$. Clearly, this does not exist for arbitrary hypersurfaces Σ. One instead picks a *polarization* $\mathcal{P}$ of $L_\Sigma^{\mathbb{C}}$, that is a subset (the 'subset of positive frequency modes') $\mathcal{P} \subset L_\Sigma^{\mathbb{C}}$ of the complexification of L_Σ such that for the symplectic complement $\mathcal{P}^\perp$, defined as the set of all $\xi \in L_\Sigma^{\mathbb{C}}$ such that $\omega_\Sigma(\xi,\eta) = 0$ for all $\eta \in \mathcal{P}$, we have $\mathcal{P}^\perp = \mathcal{P}$ (that is, $\mathcal{P}$ is Lagrangian). If $\mathcal{P}$ is a polarization, then so is $\overline{\mathcal{P}}$, where the bar denotes complex conjugation in the usual sense. $\mathcal{P}$ is called a *Kähler polarization* if $\mathcal{P} \cap \overline{\mathcal{P}} = \{0\}$. In this case, we can write $L_\Sigma^\mathbb{C} = \mathcal{P} \oplus \overline{\mathcal{P}}$.

Kähler polarizations are in one-to-one correspondence with complex structures compatible to ω_Σ. We say $J_\Sigma : L_\Sigma \to L_\Sigma$ is a complex structure compatible with ω_Σ if $J_\Sigma^2 = -\mathrm{id}_\Sigma$ and $\omega_\Sigma(J_\Sigma \cdot, J_\Sigma \cdot) = \omega_\Sigma(\cdot, \cdot)$. If such a complex structure is given, then set

$$\begin{aligned} \mathcal{P} &:= \{\phi \in L_\Sigma^\mathbb{C} | J_\Sigma \phi = -i\phi\} \\ \overline{\mathcal{P}} &:= \{\phi \in L_\Sigma^\mathbb{C} | J_\Sigma \phi = i\phi\} \end{aligned}$$

$\mathcal{P}$ and $\overline{\mathcal{P}}$ are polarizations, called the holomorphic and the antiholomorphic polarization. On the other hand, if $\mathcal{P}$ and $\overline{\mathcal{P}}$ are given Kähler polarizations, we can decompose any $\phi \in L_\Sigma^\mathbb{C}$ uniquely into $\phi = \psi + \psi'$ with $\psi \in \mathcal{P}$ and $\psi' \in \overline{\mathcal{P}}$. Then J_Σ defined by $J_\Sigma \phi := -i\psi + i\psi'$ is a complex structure compatible with ω_Σ. For the proof, see [EMRV99].

We say that J, or equivalently $\mathcal{P}$, is *positive* if $g_\Sigma(\cdot, \cdot) := 2\omega_\Sigma(\cdot, J_\Sigma \cdot)$ is a positive definite metric on L_Σ. In the following, unless otherwise stated, we shall always assume J to be a positive compatible complex structure.

If J_Σ is given, then the sesquilinear form

$$\{\phi, \eta\}_\Sigma := g_\Sigma(\phi, \eta) + 2i\omega(\phi, \eta) \quad \forall \phi, \eta \in L_\Sigma$$

turns the real vector space L_Σ into a complex Hilbert space, where multiplication with i is given by applying J_Σ. Real linear maps will lift to complex linear maps exactly if they commute with J_Σ. We have a canonical isomorphism $\pi : \mathcal{P} \to L_\Sigma$, $\pi(\psi) = \psi + \bar{\psi}$ that identifies $\mathcal{P}$ and L_Σ, the latter now regarded as a complex vector space. The inverse is explicitly given in terms of the complex structure as

$$\pi^{-1} = \frac{1}{2}(1 + iJ_\Sigma).$$

We will see later that L_Σ as a complex space can be interpreted as the one-particle Hilbert space. The state space $\mathcal{H}_\Sigma^h$ of the holomorphic quantization is now constructed as $\mathcal{H}_\Sigma^h := L^2_{hol}(L_\Sigma, d\mu_\Sigma)$[3] the space of square-integrable holomorphic functions[4] on L_Σ with respect to a certain translation invariant measure $d\mu_\Sigma$, and inner product

$$\langle \psi', \psi \rangle_\Sigma := \int_{L_\Sigma} \psi(\phi)\overline{\psi'}(\phi) \exp\left(-\frac{1}{2} g_\Sigma(\phi, \phi)\right) d\mu(\phi).$$

The amplitude ρ_M is given by

$$\rho_M(\psi) = \int_{L_{\tilde{M}}} \psi(\phi) d\nu_M(\phi)$$

where $\psi \in \mathcal{H}_{\partial M}$, $L_{\tilde{M}} \subset L_{\partial M}$ is the subspace of *global solutions* on M projected down to ∂M, and ν_M is a Gaussian probability measure determined by $\frac{1}{4} g_{\partial M}(\cdot, \cdot)$ [Oec10].

To construct the involution relating oppositely oriented manifolds explicitly we can take $\mathcal{H}_{\overline{\Sigma}}^h := L^2_{antihol}(L_\Sigma, d\mu_\Sigma)$, that is, the antiholomorphic functions. when viewed as functions on L_Σ we then have explicitly that

$$\iota_\Sigma \psi := \overline{\psi}\,. \tag{4}$$

---

[3] To deal with technical difficulties arising in the infinite dimensional case, one actually has to construct $\mathcal{H}_\Sigma^h := L^2_{hol}(\hat{L}_\Sigma, d\mu_\Sigma)$, where $\hat{L}_\Sigma$ is a certain extension of L_Σ, namely the algebraic dual of it's topological dual. The full construction of $\hat{L}_\Sigma$ and $d\mu_\Sigma$ is given in [Oec10].

[4] We say a function $f : L \to \mathbb{C}$ is holomorphic if it is continuous, bounded on every ball and holomorphic in the usual sense at every point of L.

for all $\psi \in \mathcal{H}_\Sigma$.

We will see further down, when considering the explicit construction of coherent states, that we can identify $L^2_{antihol}(L_\Sigma, d\mu_\Sigma)$ with $L^2_{hol}(L_{\overline{\Sigma}}, d\mu_\Sigma)$, where $L_{\overline{\Sigma}}$ is the same real linear space as L_Σ, but equipped with the structures $\omega_{\overline{\Sigma}} = -\omega_\Sigma$ and $J_{\overline{\Sigma}} = -J_\Sigma$.

It was shown in [Oec11] that symmetric bilinear forms Ω which define vacuum states in the Schrödinger quantization as in (3) are in one-to-one correspondence to compatible complex structures. Define $j_\Sigma : Q_\Sigma \to L_\Sigma$ to be the unique linear map such that $q_\Sigma \circ j_\Sigma = \mathrm{id}_{Q_\Sigma}$ and $j_\Sigma(Q_\Sigma) = J_\Sigma M_\Sigma$. Then

$$\Omega_\Sigma(\phi, \phi') = g_\Sigma(j_\Sigma(\phi), j_\Sigma(\phi')) - i[j_\Sigma(\phi), j_\Sigma(\phi')]_\Sigma$$

sets up the correspondence between Ω_Σ and J_Σ.

3.2.1. Coherent States

For us, coherent states will be crucially important. Fixing any element (that is, classical solution) $\xi \in L_\Sigma$, the corresponding coherent state $K_\xi \in \mathcal{H}_\Sigma$ is the holomorphic function

$$K_\xi(\phi) := \exp\left(\frac{1}{2}\{\xi, \phi\}_\Sigma\right) \qquad \forall \phi \in L_\Sigma\,. \tag{5}$$

Now if we reverse the orientation of Σ, $\Sigma \to \overline{\Sigma}$, then K_ξ gets mapped to the antiholomorphic function $\iota_\Sigma K_\xi = \overline{K_\xi}$, and comparing with the RHS of (5) we see that $\overline{K_\xi}$ is the same function as K_ξ defined with $\omega_{\overline{\Sigma}} = -\omega_\Sigma$ and $g_{\overline{\Sigma}} = g_\Sigma$ and therefore $J_{\overline{\Sigma}} = -J_\Sigma$:

$$\overline{\{\xi, \phi\}_\Sigma} = g_\Sigma(\xi, \phi) - i\omega_\Sigma(\xi, \phi) = g_{\overline{\Sigma}}(\xi, \phi) + i\omega_{\overline{\Sigma}}(\xi, \phi) = \{\xi, \phi\}_{\overline{\Sigma}}. \tag{6}$$

With respect to the inner product defined above, a coherent state has the reproducing kernel property $\langle K_\xi, \psi\rangle_\Sigma = \psi(\xi)\ \forall \psi \in \mathcal{H}_\Sigma$. One obtains for the amplitude

$$\rho_M(K_\tau) = \exp\left(\frac{1}{4}g_{\partial M}(\tau^R, \tau^R) - \frac{1}{4}g_{\partial M}(\tau^I, \tau^I) - \frac{i}{2}g_{\partial M}(\tau^R, \tau^I)\right)\,, \tag{7}$$

where $\tau = \tau^R + J_{\partial M}\tau^I$ with $\tau^R, \tau^I \in L_{\tilde{M}}$. This decomposition is always uniquely possible since $L_{\partial M} = L_{\tilde{M}} \oplus J_{\partial M} L_{\tilde{M}}$, see Proposition 4.2 in [Oec10].

There is a distinguished coherent state: The state $K_0 = 1$, the constant function with value 1 on the hypersurface Σ, which is the coherent state associated with the vector $0 \in L_\Sigma$. This state has a natural interpretation as the vacuum state on Σ. It satisfies all vacuum axioms posed in [Oec10].

If Σ decomposes into two components, $\Sigma = \Sigma_1 \cup \bar{\Sigma}_2$, then for any $\xi \in L_{\Sigma_1}$ and $\xi' \in L_{\Sigma_2}$, we have using equation (4)

$$K_{(\xi,\xi')} = K_\xi \otimes \overline{K_{\xi'}} \in \mathcal{H}_\Sigma.\,. \tag{8}$$

3.2.2. Time evolution and Unitarity

Let M be a region such that $\partial M = \Sigma_1 \cup \bar{\Sigma}_2$ is the disjoint union of two hypersurfaces. We can consider canonical projections $r_1 : L_{\tilde{M}} \to L_{\Sigma_1}$ and $r_2 : L_{\tilde{M}} \to L_{\Sigma_2}$[5]. Assume these maps are homeomorphisms. The composition $T := r_2 \circ r_1^{-1} : L_{\Sigma_1} \to L_{\Sigma_2}$ evolves solutions at Σ_1 into solutions at Σ_2, hence it can be seen as a generalized notion of time evolution. We call T unitary if

$$J_{\Sigma_2} \circ T = T \circ J_{\Sigma_1}\,. \tag{9}$$

---

[5] For $\phi \in L_{\tilde{M}}$ a global solution on M, $r_1(\phi)$ is the germ of ϕ at Σ_1, in other words $r_1(\phi)$ is obtained by forgetting ϕ everywhere but in a small neighborhood around Σ_1. Same for r_2.

In [Oec10] it was shown that if T is unitary, then there is a unitary map $U : \mathcal{H}_{\Sigma_1} \to \mathcal{H}_{\Sigma_2}$, $\Psi \mapsto \Psi \circ T^{-1}$. In particular we have

$$UK_{\Sigma_1,\xi} = K_{\Sigma_2,T\xi}.$$

Obviously $T0 = 0$ since T is linear. So the vacuum state on Σ_1 will evolve into the vacuum state on Σ_2, allowing us to construct a vacuum state $K_{(0,0)}$ on $\Sigma_1 \cup \Sigma_2$ which is compatible with the dynamics in the aforementioned sense. Since we can extend this construction to any hypersurface Σ, there will be a global vacuum state compatible with the dynamics if T is unitary.

Using the unitary map U the amplitude map can be expressed using the inner product in the form

$$\rho_M(\Psi_1 \otimes \iota_{\Sigma_2}\Psi_2) = \rho_M(\Psi_1 \otimes \overline{\Psi}_2) = \langle U^{-1}\Psi_2, \Psi_1\rangle_{\Sigma_1} \tag{10}$$

for all $\Psi_1 \in \mathcal{H}_{\Sigma_1}$ and $\Psi_2 \in \mathcal{H}_{\Sigma_2}$.

A generalization of the above definition of unitarity can be given that avoids the assumption of the existence of the evolution map T. Using the reproducing kernel property of the coherent states [Oec10] we can define a map $U^{-1} : \mathcal{H}_{\Sigma_2} \to \mathcal{H}_{\Sigma_1}$ such that (10) is fulfilled as

$$(U^{-1}\Psi)(\phi) := \rho_M(K_{\Sigma_1,\phi} \otimes \overline{\Psi})\,. \tag{11}$$

for all $\Psi \in \mathcal{H}_{\Sigma_2}$. We can extend the subspace $L_{\tilde{M}}$ to the complexification $L^{\mathbb{C}}_{\tilde{M}} \subset L^{\mathbb{C}}_{\partial_M}$, such that for $\xi \in L^{\mathbb{C}}_{\tilde{M}}$ we have $\mathfrak{Re}(\xi), \mathfrak{Im}(\xi) \in L_{\tilde{M}}$. We then have that if (9) holds every element $\xi \in L^{\mathbb{C}}_{\tilde{M}}$ that projects to an $-i$ eigenfunction of J_{Σ_1}, meaning $J_{\Sigma_1} r_1(\xi) = -ir_1(\xi)$, projects to an $-i$ eigenfunction on Σ_2, $J_{\Sigma_2} r_2(\xi) = -ir_2(\xi)$, and vice versa. Reversely we will say that a complex structure is "unitary" exactly if this condition is satisfied.

If furthermore the maps r_1 and r_2 are homeomorphisms we can identify $L^{\mathbb{C}}_{\tilde{M}}$ with $L^{\mathbb{C}}_{\Sigma_1}$ and $L^{\mathbb{C}}_{\Sigma_2}$. If the complex structure then is unitary in the above sense we find that $J_{\Sigma_2} \circ T = T \circ J_{\Sigma_1}$ since $L^{\mathbb{C}}_{\tilde{M}}$ is the set of all $(\xi, T\xi) \in L^{\mathbb{C}}_{\partial M}$.

3.2.3. Ladder Operators and one particle states

In this section, we work out the Fock space structure of $\mathcal{H}^h_\Sigma$, the state space of holomorphic quantization. Creation and annihilation operators are constructed, and the one-particle Hilbert space is constructed.

For an element $\xi \in L_\Sigma$ define the object $p_\xi \in \mathcal{H}_\Sigma$ by

$$p_\xi(\phi) = \frac{1}{\sqrt{2}}\{\xi, \phi\}_\Sigma. \tag{12}$$

We call p_ξ the normalized one-particle state corresponding to $\xi \in L_\Sigma$, for reasons to be clarified later on. In [Oec12b], the actions of creation and annihilation operators on $\mathcal{H}^h_\Sigma$ have been worked out:

$$(a^\dagger_\xi\psi)(\phi) = p_\xi(\phi)\psi(\phi) = \frac{1}{\sqrt{2}}\{\xi, \phi\}_\Sigma\psi(\phi) \tag{13}$$

$$(a_\xi\psi)(\phi) = \langle K_\phi, a_\xi\psi\rangle_\Sigma = \langle a^\dagger_\xi K_\phi, \psi\rangle_\Sigma \tag{14}$$

Here, $\xi \in L_\Sigma$, and we may say that $a^\dagger_\xi$ and a_ξ create and annihilate 'a particle of type ξ', respectively. The action of a_ξ on coherent states is particularly simple:

$$a_\xi K_\phi = \frac{1}{\sqrt{2}}\{\phi, \xi\}_\Sigma K_\phi$$

In particular, all annihilation operators annihilate the vacuum K_0. These operators satisfy the usual commutation relations

$$[a_\xi, a^\dagger_\eta] = \{\eta, \xi\}_\Sigma, \qquad [a_\xi, a_\eta] = 0, \qquad [a^\dagger_\xi, a^\dagger_\eta] = 0. \tag{15}$$

The creation and annihilation operators together with the vacuum state determine a Fock structure on $\mathcal{H}^h_\Sigma$. We wish to determine the one-particle sector of this Fock structure. This will be done by defining the number operator. For that purpose, we first define an orthonormal basis in $\mathcal{P}$:

Let $\pi : \mathcal{P} \to L_\Sigma$ be the canonical linear isomorphism, with $\pi(\phi) = \phi + \bar{\phi}$. We use this isomorphism to define an inner product on $\mathcal{P}$:

$$\{\cdot, \cdot\}_\mathcal{P} := \{\pi(\cdot), \pi(\cdot)\}_\Sigma.$$

Using linearity of π and the fact that $\{\cdot, \cdot\}_\Sigma$ is an inner product on L_Σ, it is immediate to show that $\{\cdot, \cdot\}_\mathcal{P}$ is an inner product on $\mathcal{P}$. Now, let $\mathcal{B}_\mathcal{P} \subset \mathcal{P}$ be an orthonormal basis of $\mathcal{P}$ with respect to this inner product. We shall write $\mathcal{B}_\mathcal{P} = \{e^+_\xi\}_{\xi \in \mathcal{I}}$ for a suitable index set $\mathcal{I}$.

Now we define the number operator in that basis $\mathcal{B}_\mathcal{P}$

$$N_\Sigma = \sum_{\xi \in \mathcal{B}_\mathcal{P}} a^\dagger_{\pi(\xi)} a_{\pi(\xi)} \equiv \sum_{\xi \in \mathcal{B}_\mathcal{P}} a^\dagger_\xi a_\xi \tag{16}$$

in the usual sense, where, using π to identify $\mathcal{P}$ and complex L_Σ, we introduced the shorthand notation $a_{\pi(\xi)} = a_\xi$.

Write $|0\rangle = K_0$. Using the commutation relations it is then easy to see that for $\eta \in \mathcal{B}_\mathcal{P}$,

$$N_\Sigma a^\dagger_\eta |0\rangle = \sum_\xi a^\dagger_\xi a_\xi a^\dagger_\eta |0\rangle = \sum_\xi a^\dagger_\xi [a_\xi, a^\dagger_\eta] |0\rangle = \sum_\xi a^\dagger_\xi \{\eta, \xi\}_\Sigma |0\rangle = a^\dagger_\eta |0\rangle.$$

The linear space spanned by the vectors $a^\dagger_\eta|0\rangle$ for $\eta \in \mathcal{B}_\mathcal{P}$ is thus the eigenspace of N_Σ corresponding to the eigenvalue 1. Using (13), it is easy to see that

$$a^\dagger_{c_1\xi_1 + \ldots c_n\xi_n} = c_1 a^\dagger_{\xi_1} + \ldots c_n a^\dagger_{\xi_n}, \quad c_i \in \mathbb{C}, \xi_i \in \mathcal{B}_\mathcal{P}.$$

This means the set

$$\left\{a^\dagger_\eta|0\rangle, \eta \in \mathcal{P}\right\} = \left\{\frac{1}{\sqrt{2}}\{\eta, \cdot\}, \eta \in L_\Sigma\right\} = \{p_\eta, \eta \in L_\Sigma\} =: L^*_\Sigma \simeq L_\Sigma$$

is the one-particle sector of $\mathcal{H}^h_\Sigma$, and we were indeed justified to call $p_\xi(\phi) = \frac{1}{\sqrt{2}}\{\xi, \phi\}_\Sigma$ a one-particle-state.

4. Choosing a complex structure

As we saw in section 3.2, the remaining quantization ambiguity is the choice of a suitable positive complex structure J which is compatible with ω. In this section we study a number of physically motivated conditions that allow us to select such a complex structure. For technical simplicity, our field theory of interest will be ordinary scalar field theory in Minkowski spacetime. The analysis is divided into two parts: First we study quantization in regions bounded by spacelike hypersurfaces, which is well understood. The right condition for the selection of J turns out to be the *Ashtekar-Magnon condition* [AM75]. We then study quantization in regions bounded by timelike hypersurfaces using GBQFT. It turns out that, in comparison to the spacelike case, solutions to the classical field equations arise which have the behavior of evanescent waves. These solutions cause a number of problems which will be discussed below. We will give two possible generalizations of the Ashtekar-Magnon condition for timelike hypersurfaces, which both work for all solutions except the aforementioned evanescent waves.

4.1. Spacelike hypersurfaces - The Ashtekar-Magnon energy condition

In [AM75] Ashtekar and Magnon address the question of finding a positive compatible complex structure within the framework of geometric quantization. They show that if an *energy condition* is imposed, the complex structure is uniquely determined. The energy condition amounts to asking that the expectation value of the single particle Hamiltonian operator be the same as the classical energy of the field. It replaces the standard 'positive frequency modes' condition which relies on the presence of a global timelike Killing vector field and works for a wider class of spacetimes which admit a foliation into Cauchy hypersurfaces. We present their result here as the starting point of our discussion about complex structures. Some notation first: With the definitions of section 3.1, there is a map F_Σ from $L_\Sigma = N_\Sigma \oplus M_\Sigma$ to the phase space $\hat{L}_\Sigma = C_0^\infty(\Sigma, \mathbb{R}) \oplus C_0^\infty(\Sigma, \mathbb{R})$, given by

$$F_\Sigma \phi = (\phi|_\Sigma, (\sqrt{h} n^\mu \nabla_\mu \phi)|_\Sigma)\,, \tag{17}$$

The phase space can be seen as the space of initial data. If Σ is a Cauchy hypersurface, then by the Cauchy property F_Σ is an isomorphism, and $L_\Sigma \simeq \hat{L}_\Sigma$. The structures on L_Σ then all have obvious counterparts on $\hat{L}_\Sigma$. In addition, an operator J_Σ on L_Σ corresponds uniquely to an operator on the phase space. Ashtekar and Magnon use this phase space representation to prove the following theorem:

Theorem 1 (Ashtekar-Magnon) *Let (M, g) be a globally hyperbolic spacetime*[6] *with Cauchy surface Σ, and L_Σ as above. Then there exists a unique complex structure J on L_Σ with the following properties :*

1. *J is positive and compatible with ω, that is $\omega(\cdot, J(\cdot)) \geq 0$ and $\omega(J(\cdot), J(\cdot)) = \omega(\cdot, \cdot)$.*
2. *The expectation value of the one-particle Hamiltonian operator equals the classical energy* [7]*:*

$$\{\phi, H\phi\}_\Sigma = -2 \int_\Sigma d\sigma_\mu T^{\mu\nu}(\phi) t_\nu \qquad \forall \phi \in L_\Sigma \tag{18}$$

where $H\phi = -J\dot{\phi}$, and $\dot{\phi}$ is the solution [8]

$$\dot{\phi} = F_\Sigma^{-1} \left(t^\mu \nabla_\mu \phi|_\Sigma, t^\mu \nabla_\mu \sqrt{h} n^\nu \nabla_\nu \phi|_\Sigma \right)$$

and t is the vector field corresponding to the hypersurface orthogonal foliation by Cauchy hypersurfaces Σ_t in a neighborhood of Σ.

In [FD99] it was shown that the real part of (18)

$$\omega_\Sigma(\phi, \dot{\phi}) = -2 \int_\Sigma d\sigma_\mu T^{\mu\nu}(\phi) t_\nu$$

which is independent of J is actually always true, so that the true content of the energy condition is it's imaginary part:

$$\omega_\Sigma(\phi, H\phi) = 0 \qquad \forall \phi \in L_\Sigma. \tag{19}$$

Furthermore, in [FD99] the action of the complex structure found by Ashtekar and Magnon on the phase space is worked out as

---

[6] Our signature convention for the metric is $(+ - - -)$.

[7] The additional factor of (-2) in comparison with the expression by Ashtekar and Magnon is due to our different definition of the symplectic form ω and $\{\cdot, \cdot\}$.

[8] The dot does not refer to a time derivative in the usual sense! However, if there is a global killing vector field t which is orthogonal to Σ, then $\dot{\phi} = \mathcal{L}_t \phi$, where $\mathcal{L}_t$ is Lie differentiation with respect to t.

$$J_\Sigma = F_\Sigma^{-1} \circ \begin{pmatrix} & \Theta^{-1/2}\frac{N}{\sqrt{h}} \\ -\frac{\sqrt{h}}{N}\Theta^{1/2} & \end{pmatrix} \circ F_\Sigma \tag{20}$$

where

$$\Theta = -(N^2 h^{ij} D_i D_j + h^{ij} N D_i (N D_j) - m^2 N^2) \tag{21}$$

and $\sqrt{h} = (\det h)^{1/2}$ with $-h$ the pullback of g to the hypersurface and $N = \sqrt{g_{00}}$ the lapse function. Using that $J_\Sigma^2 = -1$ we find

$$J_\Sigma \phi = \Theta^{-1/2} N n^\mu \nabla_\mu \phi. \tag{22}$$

To use this complex structure we will systematically excise the mode with $\Theta = 0$.

4.1.1. The canonical example: Equal-time hyperplanes

In this section we briefly review the standard construction of free scalar field theory in the GBF framework. In what follows, take admissible hypersurfaces to be oriented equal-time hypersurfaces, and admissible regions to be regions whose boundary decomposes into such hypersurfaces. The analysis in this and the next subsection was done in [Oec10].

We parametrize global solutions of the Klein-Gordon equation in the usual way via plain waves,

$$\phi(t,x) = \int \frac{\mathrm{d}^3 p}{(2\pi)^3 2p_0} \left(\phi(\vec{p}) e^{-i(p_0 x_0 - \vec{p}\cdot\vec{x})} + c.c. \right), \tag{23}$$

where $\phi : \mathbb{R}^3 \to \mathbb{C}$ is a complex function on momentum space of a type to be determined below and $p_0 := \sqrt{\vec{p}^2 + m^2}$. The symplectic form on L_{Σ_t} is

$$\begin{aligned} \omega_{\Sigma_t}(\phi_1, \phi_2) &= \tfrac{1}{2} \int \mathrm{d}^3 x \ \left(\phi_2(x_0, \vec{x}) \partial_0 \phi_1(x_0, \vec{x}) - \phi_1(x_0, \vec{x}) \partial_0 \phi_2(x_0, \vec{x}) \right)|_{\Sigma_t} \\ &= \mathfrak{Im} \left(\int \tfrac{\mathrm{d}^3 p}{(2\pi)^3 2p_0} \phi_1(\vec{p}) \overline{\phi_2(\vec{p})} \right). \end{aligned} \tag{24}$$

We chose the orientation of Σ_t to be that induced by a future pointing normal. We can invoke the AM-condition and the equations of motion to find the standard complex structure

$$J_{\Sigma_t} = \frac{\partial_0}{\sqrt{-\partial_0^2}}. \tag{25}$$

This complex structure produces a unitary time evolution. The induced metric is

$$g_{\Sigma_t}(\phi_1, \phi_2) = 2\,\mathfrak{Re} \left(\int \frac{\mathrm{d}^3 p}{(2\pi)^3 2p_0} \phi_1(\vec{p}) \overline{\phi_2(\vec{p})} \right). \tag{26}$$

and this leads to the standard complex inner product on L_{Σ_t},

$$\{\phi_1, \phi_2\}_{\Sigma_t} = 2 \int \frac{\mathrm{d}^3 \vec{p}}{(2\pi)^3 2p_0} \, \phi_1(\vec{p}) \overline{\phi_2(\vec{p})}. \tag{27}$$

This also determines the space L_{Σ_t} precisely as the space of (equivalence classes of) square-integrable complex functions $\phi : \mathbb{R}^3 \to \mathbb{C}$ with the inner product written above. The quantization thus obtained is equivalent to the usual well-known quantization of Klein-Gordon theory.

4.2. The stress-momentum condition

In this section we investigate another quantum-classical correspondence condition that we can impose for general hypersurfaces. We will see that this reproduces part of the symmetry considerations that determine the complex structure in the standard treatment.

Let Σ be a spacelike or timelike hyperplane in Minkowski space. We impose the following stress-momentum conditions

1. J is compatible with ω_Σ, that is $\omega_\Sigma(\cdot, J_\Sigma(\cdot))$ is positive definite and $\omega_\Sigma(J_\Sigma(\cdot), J_\Sigma(\cdot)) = \omega_\Sigma(\cdot, \cdot)$.
2. The expectation value of the one-particle momentum operator $\Pi := J_\Sigma t^\mu \partial_\mu$, with t^μ a constant tangent vector field to the boundary, equals the classical momentum:

$$\{\phi, \Pi\phi\}_\Sigma = -2 \int_\Sigma d\sigma_\mu T^{\mu\nu}(\phi) t_\nu \qquad \forall \phi \in L_\Sigma \, , \tag{28}$$

It is again easy to see that (28) splits into the conditions

$$\begin{aligned} \omega_\Sigma(\phi, t^\mu \partial_\mu \phi) &= \int_\Sigma d\sigma_\mu T^{\mu\nu}(\phi) t_\nu \\ \omega_\Sigma(\phi, \Pi\phi) &= 0 \end{aligned}$$

for all $\phi \in L_\Sigma$. The first condition is always fulfilled which can be proven easily integrating by parts in ω. Using Lemma 1 from [FD99] we find that the second condition is the statement that Π is a symmetric operator with respect to the inner product $\{\cdot, \cdot\}_\Sigma$. With Lemma 2 follows then that $[J_\Sigma, t^\mu \partial_\mu] = 0$. We obtain that the condition in (28) is equivalent to the invariance of the complex structure under translation in t direction. Since this is a symmetry of our classical configuration it is a completely natural condition.

We conclude that a complex structure that satisfies (28) leaves the eigenspaces of $t^\mu \partial_\mu$ invariant and there exists a complete set of eigenfunctions of J_Σ where every element consists of linear combinations of eigenfunctions of $t^\mu \partial_\mu$ to the same eigenvalue.

4.3. Timelike hyperplanes

In this section we consider timelike hyperplanes in $(3+1)$-Minkowski spacetime. We take $g = \mathrm{diag}(1, -1, -1, -1)$. Admissible hypersurfaces are all oriented hyperplanes perpendicular to the x_1-axis and their finite disjoint unions. Admissible regular regions are all finite closed intervals on the x_1-axis extended in the other coordinate directions and their finite disjoint unions. We assume the hyperplanes to be oriented in negative x_1 direction.

The AM-condition does not apply directly to this scenario, but we can try to adapt it by choosing different components of the energy-momentum tensor and forcing them to coincide with their classical counterparts in the one particle sector.

4.3.1. The pressure condition

In section 4.1 we saw that the AM-condition for the directions of the momentum in the hyperplane reproduce the symmetry considerations familiar from the standard treatment in Minkowski space for all directions tangent to the hyperplane. In order to fully determine the complex structure we now need to consider the orthogonal direction.

Here we cannot apply a direct analogue of the theorem of Ashtekar and Magnon. We don't have Cauchy hypersurfaces, and there is no notion of phase space immediately available. Therefore there is no sensible definition of a Hamiltonian operator at hand. Nevertheless, a few statements can be made. Firstly, for one admissible hypersurface Σ, let n be the unit vector field orthogonal to Σ. Without loss of generality we require n to point in the positive x_1 direction. Due to the symmetries of Minkowski space n is non-vanishing and killing everywhere, which allows us to define $\dot{\phi} = \mathcal{L}_n \phi = n^\mu \partial_\mu \phi$ even in the absence of a phase space. With this in mind, we write in analogy to the real part of (18):

$$\mathfrak{Re}\{\phi, H\phi\}_\Sigma = 2\omega_\Sigma(\phi, n^\mu \partial_\mu \phi) = -2 \int_\Sigma d\sigma_\mu T^{\mu\nu}(\phi) n_\nu. \qquad \forall \phi \in L_\Sigma \tag{29}$$

for some operator H. We give a proof in appendix A.1 that the second equality is always satisfied, and the first equality defines H: Using the definition of $\{\cdot,\cdot\}$, we see that the operator H must satisfy $J_\Sigma H = n^\mu \partial_\mu =: D^\perp$ or $H = -J_\Sigma D^\perp$. Now of course the RHS does no longer have the interpretation of a classical energy. In our case, we might interpret it as a classical momentum corresponding to the x_1-direction.

Now we may ask the question 'is there a complex structure J which is

(1) positive and compatible with ω_Σ

(2) satisfies the imaginary part of (18), that is $\omega_\Sigma(\phi, H\phi) = 0$ for $H = -J_\Sigma D^\perp$?'

We call this condition 'pressure condition' since the RHS of (29) suggests that we are looking for a quantum observable corresponding to the T^{11}-component of the energy-momentum tensor, which has the interpretation of a pressure.

Now we are tempted by the analysis of the spatial case to set J in analogy to (22)

$$J_\Sigma \phi = -\Theta^{-1/2} N n^\mu \nabla_\mu \phi \tag{30}$$

with

$$\Theta = (N^2 h^{ij} D_i D_j + h^{ij} N (D_i N D_j) - m^2 N^2).$$

Here Θ is defined with the opposite sign with respect to (21) for convenience. Now n^μ is the vector field orthogonal to Σ, $-h$ is the pullback of the metric to Σ and $N = \sqrt{g_{11}} = 1$. So Θ reduces to

$$\Theta = h^{ij} \partial_i \partial_j - m^2 = -\partial_0^2 + \partial_2^2 + \partial_3^2 - m^2 = -\partial_1^2$$

where the last equality follows from the equations of motion. The problem with this operator is that it does not have a positive spectrum, so we cannot construct $\Theta^{1/2}$. This can be seen directly from the Fourier transform $\tilde{\Theta}$ of Θ which is

$$\tilde{\Theta} = p_0^2 - p_2^2 - p_3^2 - m^2,$$

and by the equations of motion this is equal to p_1^2. The positive part of the spectrum of Θ are those values of (p_0, p_2, p_3) for which $\tilde{\Theta} > 0$, which implies that p_1 is real. We call solutions which are eigenvectors of Θ corresponding to positive eigenvalues *propagating waves.* They span a subspace $L_\Sigma^p \subset L_\Sigma$, and on that subspace Θ is positive self-adjoint which allows us to define $\Theta^{1/2}$, then J can indeed be defined by (30) on that subspace, which yields

$$J_\Sigma^p = -\frac{\partial_1}{\sqrt{-\partial_1^2}}\,. \tag{31}$$

An interesting fact about this complex structure is that among all complex structures which are of the form $\frac{\partial_\eta}{\sqrt{-\partial_\eta^2}}$ for some vector field η (which is the form we found in section 4.1.1), J_Σ^p is the only one which actually satisfies condition (1), i.e. is compatible with ω [9]. The proof for this statement can be found in the appendix A.2. Thus we seem to have little choice in defining J_Σ, even if condition (2) is left aside completely.

---

[9] It might be interesting to note that this is not true in the case of spacelike hyperplanes, where the allowed complex structures correspond to all vector fields which are inside the forward light cone on every point in Σ

The negative part of the spectrum of Θ are those values of (p_0, p_2, p_3) for which $\tilde{\Theta} < 0$, which implies that p_1 is imaginary. We call solutions which are eigenvectors of Θ corresponding to negative eigenvalues *evanescent waves*. These are additional solutions which decay or grow exponentially as one moves away from the hypersurface. They are thus well-behaved near the hypersurface, but not globally. They span a subspace $L^e_\Sigma \subset L_\Sigma$, and $L^e_\Sigma \cup L^p_\Sigma = L_\Sigma$. One could replace Θ by $-\Theta$ on L^e_Σ to obtain a positive self-adjoint operator. However, we then have to find an operator L such that $J^e_\Sigma = L(-\Theta)^{-1/2}$ squares to minus one. One possible choice for the timelike hyperplane Σ is $L = P_\Sigma \partial_1$, where P_Σ is reflection along Σ, such that $P_\Sigma f(x_{||}, x_\perp) = f(x_{||}, -x_\perp)$ if $x_\perp$ is the coordinate orthogonal to Σ. We have that $P_\Sigma^2 = 1$ and P_Σ anti-commutes with ∂_1, so that we can write

$$J^e_\Sigma = -\frac{P_\Sigma \partial_1}{\sqrt{\partial_1^2}}.$$

One can check that this complex structure is indeed compatible with ω on L^e_Σ.

Now what about condition (2)? We have two different complex structures on our two subspaces L^p_Σ and L^e_Σ, and by $H = -J_\Sigma D^\perp$ we will also have two different 'Hamiltonians'. First consider propagating waves. We can write $J^p_\Sigma = -D^\perp \Theta^{-1/2}$, so to satisfy $J^p_\Sigma H_p = D^\perp$ we must have $H_p = -\Theta^{1/2}$. This is a symmetric operator, and as such indeed satisfies $\omega_\Sigma(\phi, H_p\phi) = 0$: On the one hand we have $\omega_\Sigma(\phi, H_p\phi) = -\omega_\Sigma(H_p\phi, \phi)$ because ω_Σ is antisymmetric, on the other hand $\omega_\Sigma(\phi, H_p\phi) = \omega_\Sigma(H_p\phi, \phi)$ since H_p is symmetric.

Now consider evanescent waves. We have $J^e_\Sigma = -(-\Theta)^{-1/2} P_\Sigma \partial_1$, and

$$H_e = -J^e_\Sigma D^\perp = (-\Theta)^{-1/2} P_\Sigma \partial_1 P_\Sigma^2 \partial_1 = -(-\Theta)^{-1/2} (P_\Sigma \partial_1)^2 P_\Sigma = (-\Theta)^{1/2} P_\Sigma.$$

Now the operator $(-\Theta)^{1/2}$ is symmetric and thus satisfies

$$\omega_\Sigma\left(\phi, (-\Theta)^{1/2}\psi\right) = \omega_\Sigma\left((-\Theta)^{1/2}\phi, \psi\right).$$

The operator P_Σ, however is antisymmetric which can be seen from $P_\Sigma^2 = 1$ and the observation that

$$\omega_\Sigma\left(P_\Sigma\phi, P_\Sigma\psi\right) = -\omega_\Sigma\left(\phi, \psi\right). \tag{32}$$

As a consequence H_e will be antisymmetric which can be seen by using the explicit expression for ω_Σ on the timelike hyperplane

$$\omega_{\Sigma_\xi}(\eta_1, \eta_2) = \frac{1}{2} \int \mathrm{d}x_0 \, \mathrm{d}^2\tilde{x} \;\; \left(\eta_1(t, x_1, \tilde{x}) \partial_1 \eta_2(t, x_1, \tilde{x}) - \eta_2(t, x_1, \tilde{x}) \partial_1 \eta_1(t, x_1, \tilde{x})\right)|_{\Sigma_\xi}. \tag{33}$$

Since ω_Σ is non-degenerate there is a ψ' for every ϕ such that $\omega_\Sigma(\phi, \psi') \neq 0$. Now define ψ such that $\psi' = H_e\psi$ which is always possible since H_e is invertible, then $\omega_\Sigma(\phi+\psi, H_e(\phi+\psi)) = 2\omega_\Sigma(\phi, H_e\psi) \neq 0$. Hence, $\omega_\Sigma(\phi, H_e\phi)$ will not be zero in general. Thus the Ashtekar-Magnon condition fails to hold for evanescent waves. Essentially we have no good pressure operator for evanescent waves available.

In the following we use the complex structure J^e_Σ we derived above and J^p_Σ we defined in analogy to write explicit relations for metric and inner product on timelike hypersurfaces in Minkowski space. They coincide with those presented in [Oec10]. There the author gives a parameterization of L_{Σ_ξ} for a timelike hyperplane Σ_ξ with tangent ∂_0, ∂_2 and ∂_3 at $x_1 = \xi$ into eigenfunctions of $J_{\Sigma_\xi} = J^p_{\Sigma_\xi}|_{L^p_{\Sigma_\xi}} + J^e_{\Sigma_\xi}|_{L^e_{\Sigma_\xi}}$ via a complex function $\eta : \mathbb{R} \times \mathbb{R}^2 \to \mathbb{C}$ as:

$$\eta(x_0, x_1, \tilde{x}) = \int \frac{\mathrm{d}p_0 \, \mathrm{d}^2\tilde{p}}{(2\pi)^3 2p_1} \left(\eta(p_0, \tilde{p}) f(p_0, \tilde{p}, x_1) e^{-i(p_0 x_0 - \tilde{p}\tilde{x})} + c.c.\right), \tag{34}$$

where $\tilde{x} := (x_2, x_3)$, $\tilde{p} := (p_2, p_3)$, $p_1 := \sqrt{|p_0^2 - \tilde{p}^2 - m^2|}$ and

$$f(p_0, \tilde{p}, x_1) := \begin{cases} e^{ip_1(x_1 - \xi)} = \cos(p_1(x_1 - \xi)) + i\sin(p_1(x_1 - \xi)) & \text{if } p_0^2 - \tilde{p}^2 - m^2 > 0 \\ \cosh(p_1(x_1 - \xi)) + i\sinh(p_1(x_1 - \xi)) & \text{if } p_0^2 - \tilde{p}^2 - m^2 < 0. \end{cases} \tag{35}$$

The symplectic form in (33) on the space of solutions L_{Σ_ξ} associated to the hyperplane Σ_ξ with orientation pointing in positive x_1 direction turns out to be

$$\omega_{\Sigma_\xi}(\eta_1,\eta_2) = \mathfrak{Im}\left(\int \frac{\mathrm{d}p_0\,\mathrm{d}^2\tilde{p}}{(2\pi)^3 2p_1}\eta_1(p_0,\tilde{p})\overline{\eta_2(p_0,\tilde{p})}\right). \tag{36}$$

It can be seen that J^e and J^p together act on η as defined in (34) just by multiplication:

$$(J(\eta))(p_0,\tilde{p}) = -i\eta(p_0,\tilde{p}).$$

The resulting metric is,

$$g_{\Sigma_\xi}(\eta_1,\eta_2) = 2\mathfrak{Re}\left(\int \frac{\mathrm{d}p_0\,\mathrm{d}^2\tilde{p}}{(2\pi)^3 2p_1}\eta_1(p,\tilde{p})\overline{\eta_2(p_0,\tilde{p})}\right). \tag{37}$$

The complex inner product is then given by

$$\{\eta_1,\eta_2\}_{\Sigma_\xi} = 2\int \frac{\mathrm{d}p_0\,\mathrm{d}^2\tilde{p}}{(2\pi)^3 2p_1}\,\eta_1(p_0,\tilde{p})\overline{\eta_2(p_0,\tilde{p})}. \tag{38}$$

The space L_{Σ_ξ} is thus precisely the space of (equivalence classes of) complex valued square-integrable functions $\mathbb{R}\times\mathbb{R}^2\to\mathbb{C}$ with the above inner product.

4.3.2. Physical interpretation

We have seen that, for propagating waves the pressure condition can indeed determine the complex structure in the case of timelike hyperplanes. It is instructive to explicitly contrast the different roles the complex structures play in this case though.

Recall that we can give a compatible, though not necessarily positive, complex structure by giving a polarization, that is a subspace of $L^{\mathbb{C}}_{\Sigma}$. Let us first consider the case of a spacelike hyperplane Σ in Minkowski space. Imposing the stress-momentum condition of section 4.2 for all tangents to Σ we can split $L^{\mathbb{C}}_{\Sigma}$ into momentum eigenspaces that are not mixed by the complex structure. Thus, specializing to $1+1$ dimensions for simplicity, the problem of specifying a polarization reduces to the problem of giving a two dimensional subspace of the space spanned by the four modes $e(\pm p_0,\pm p_1)\in L^{\mathbb{C}}_{\Sigma}$ where $e(p_0,p_1)=e^{-i(p_0x_0-p_1x_1)}$ corresponds to the wave vector $p=(p_0,p_1)$. The usual choice coinciding with the complex structure specified by the AM-condition we reviewed in section 4.1 is to take the positive frequency modes $e(+p_0,\pm p_1)$.

Now considering a region $\Sigma\times[t_i,t_f]$ bounded by two spacelike hyperplanes Σ_i and Σ_f we have that Σ_f has the same orientation as the reference hyperplane Σ but Σ_i has orientation opposite to that of Σ. This is the left side of figure 1. As discussed above changing orientation is simply complex conjugation, and we have $\overline{e(p_0,p_1)}=e(-p_0,-p_1)$. Thus writing $e_i\in L^{\mathbb{C}}_{\Sigma_i}$ and $e_f\in L^{\mathbb{C}}_{\Sigma_f}$ we obtain from the standard choice of picking positive frequency modes $e_f(p_0,\pm p_1)$ and $e_i(-p_0,\mp p_1)$. These are exactly the modes for which the wave vector is pointing out of the spacetime region.

Now consider the region $\Sigma\times[x_l,x_r]$ bounded by the timelike hyperplanes Σ_l and Σ_r with modes $e_l\in L^{\mathbb{C}}_{\Sigma_l}$ and $e_r\in L^{\mathbb{C}}_{\Sigma_r}$ where we again imposed the stress-momentum conditions for all tangents to Σ_l and Σ_r. For propagating waves the complex structure derived in section 4.3.1 selects the left pointing modes $e_l(p_0,-p_1)$ and $e_l(-p_0,-p_1)$ in the left hypersurface and the right pointing modes $e_r(p_0,p_1)$ and $e_r(-p_0,p_1)$ on the right. This is indicated on the right side of figure 1. Thus we see that this complex structure again selects the modes pointing out of the spacetime region under consideration. This is thus the covariant formulation of the intuition to select positive frequency modes. We can also immediately see why this intuition becomes meaningless for evanescent waves, for which the wavevector becomes complex.

Note that the same modes are still present in the case of timelike hyperplanes as in the case of spacelike hyperplanes, however, the corresponding one particle states and even the boundary vacuum



Figure 1: The modes selected (solid arrows) and discarded (dashed arrows) by the complex structure. The orthogonal arrows indicate the hypersurface orientation. The future pointing modes on $\overline{\Sigma}_i$ become past pointing on Σ_i. Thus on Σ_i and Σ_f we have only outward pointing modes.

states must now be interpreted in a way that is less obvious than in the case of spacelike hyperplanes. To illustrate this, consider a scenario with a source inducing a transition from the vacuum to a one particle state corresponding to an arbitrary mode, as we will study in detail in section 5.3. In the case of spacelike hyperplanes, the condition that there is a vacuum in the past, that is absence of any particles is simply given by selecting the boundary vacuum state on Σ_i and leaving the states on Σ_f completely free. This has the effect of ruling out the modes $e_i(-p_0, \mp p_1)$.

In the case of timelike hypersurfaces this has to be expressed differently. Choosing the vacuum state on either side would clearly lead to an incorrect selection of modes. In particular we want to choose the modes $e_l(p_0, -p_1)$ and $e_r(p_0, p_1)$. Now consider the symmetric operator $\partial_0 \partial_\eta$, with ∂_η the normal derivative to the surface our mode is associated to. Selecting the modes corresponding to positive eigenvalues of this operator within the polarization indeed selects the correct modes.

We suggest interpreting the condition that $\partial_0 \partial_\eta$ be positive as the condition that there is a positive energy flux through the surface under consideration. This is based on the physical idea to replace the condition that we have a vacuum in the past with the notion that there should be no energy flowing into the spacetime region under consideration.

Consider the right hypersurface. We can see easily that for the solutions $\phi^+_{p_0} = ce(p_0, p_1) + \overline{c}e(-p_0, -p_1)$ for which $\partial_0 \partial_1$ is positive, the classical energy momentum tensor is also everywhere positive,

$$T^{01}(\phi^+_{p_0})(x) > 0,$$

as

$$T^{01}(\phi^+_{p_0})(x) = -(\partial_0 \phi^+_{p_0})(\partial_1 \phi^+_{p_0})(x) = p_0 p_1 {(\phi^+_{p_0})}^2(x)$$

While this is sufficient for our purposes it is necessary to generalize this condition to less symmetric situations and to the full Hilbert space of the theory, when for example, considering higher order perturbation theory. We leave that for future work.

To summarize, for propagating modes and the spacetime regions considered here we can replace the notion of positive frequency modes with the notion of modes pointing out of the spacetime region under consideration, and the creation of modes out of the vacuum by a source, can be described by restricting the state space to the modes with outward flowing energy.

5. The Unruh-DeWitt detector

As we saw in the preceding section the usual abstract arguments only determine the complex structure for propagating waves. In order to investigate the choice of complex structure for evanescent waves we will now work out how to couple a detector to a quantum field theory in the general boundary formalism. Using this model and certain symmetry conditions we recover the result of the AM like condition for propagating waves derived in the foregoing section. Then we apply this setup to a physical situation where evanescent waves cause excitations. We will see that the choice of complex structure for evanescent waves depends on the physical setup outside of the region of interest, and thus no general arguments can determine a unique complex structure for these modes.

5.1. Perturbation theory in GBQFT

From [CO08a] we find that the amplitude for two fields ϕ_1 and ϕ_2 interacting via a potential $\mathcal{V}(\phi_1, \phi_2)$ in GBQFT can be generated perturbatively from the amplitude of the affine field theory

$$S_\mu[\phi] = S_0[\phi] + \int dV(x)\mu(x)\phi(x)\,.$$

We define the term of perturbative order n as

$$\rho^S_{n,V}(K_{\xi_1} \otimes K_{\xi_2}) := \frac{1}{n!}\left(\int dV(x)\mathcal{V}\left(-i\frac{\delta}{\delta\mu_1(x)}, -i\frac{\delta}{\delta\mu_2(x)}\right)\right)^n \rho^S_{\mu_1}(K_{\xi_1})\rho^S_{\mu_2}(K_{\xi_2})\Big|_{\mu_1,\mu_2=0}\,. \tag{39}$$

We remark that the amplitude $\rho^S_\mu(K_\xi)$, the amplitude associated to the action S_μ, can equivalently be seen as the operator amplitude for the Weyl observable $W := e^{iD}$ with $D(\phi) = \int dV(x)\,\mu(x)\phi(x)$ [Oec12c]. An expression for this amplitude was derived in [Oec12c]. We have

$$\begin{aligned}\rho^S_\mu(K_\xi) &= \rho_M(K_\xi)W(\hat{\xi})e^{\frac{i}{2}D(\eta_D)-\frac{1}{2}g_{\partial M}(\eta_D,\eta_D)}\\ &= \rho_M(K_\xi)W(\hat{\xi})e^{\frac{i}{2}D((1-iJ_{\partial M})\eta_D)}\end{aligned} \tag{40}$$

where $\hat{\xi} = \xi^R - i\xi^I$ with $\xi = \xi^R + J_{\partial M}\xi^I$ and $\xi^R, \xi^I \in L_{\tilde{M}}$ as in section 3.2.1 and $\eta_D \in L_{\partial M}$ is uniquely defined by the condition that $D(\zeta) = 2\omega_{\partial M}(\zeta, \eta_D)$ for all $\zeta \in L_M$. The amplitude in equation (39) is the analog of the scattering matrix of n-th order in standard QFT. In GBQFT probabilities are always given as conditional probabilities [Oec08]. We obtain that the probability to find the interacting system in a state in the closed subset $\mathcal{J} \subseteq \mathcal{H}_{\partial M}$ knowing that it can only be in a state in the closed subset $\mathcal{I} \subseteq \mathcal{H}_{\partial M}$ that contains $\mathcal{J}$ is given as

$$P(\mathcal{J}|\mathcal{I}) = \frac{||\sum_n \rho^S_{n,V} \circ P_{\mathcal{J}}||^2}{||\sum_n \rho^S_{n,V} \circ P_{\mathcal{I}}||^2} \tag{41}$$

where the norm $||\cdot||$ is with respect to the inner product of the dual Hilbert space $\mathcal{H}^*_{\partial M}$, and $P_{\mathcal{J}}$ and $P_{\mathcal{I}}$ are the orthogonal projections to $\mathcal{J}$ and $\mathcal{I}$ respectively. Starting from an orthonormal basis in $\mathcal{J}$ which we extend first to an orthonormal basis of $\mathcal{I}$ we arrive after another extension at an orthonormal basis $\{\xi_i\}_{i\in\mathbb{N}}$ of $\mathcal{H}_{\partial M}$. Using Riesz' theorem we find that the element of $\psi_{\rho,\mathcal{J}} \in \mathcal{H}_{\partial M}$ that corresponds to the dual element $\sum_n \rho^S_{n,V} \circ P_{\mathcal{J}}$ via the inner product on $\mathcal{H}_{\partial M}$ is given as

$$\psi_{\rho,\mathcal{J}} = \sum_{\xi_i\in\mathcal{J}} \overline{\sum_n \rho^S_{n,V}(\xi_i)}\xi_i \tag{42}$$

and analogously for $\sum_n \rho^S_{n,V} \circ P_{\mathcal{I}}$. The norm of $\sum_n \rho^S_{n,V} \circ P_{\mathcal{J}}$ in $\mathcal{H}^*_{\partial M}$ can then be written as the norm of $\psi_{\rho,\mathcal{J}}$ in $\mathcal{H}_{\partial M}$ and we can rewrite the expression in equation (41) as

$$P(\mathcal{J}|\mathcal{I}) = \frac{\sum_{\xi_j \in \mathcal{J}} |\sum_n \rho^S_{n,V}(\xi_j)|^2}{\sum_{\xi_i \in \mathcal{I}} |\sum_n \rho^S_{n,V}(\xi_i)|^2} \,. \tag{43}$$

The probability for the system to be found in the normalized one-particle state $\psi \subset \mathcal{I}$ is then given in first order in the interaction

$$P(\psi|\mathcal{I}) = \frac{|\rho^S_{1,V}(\psi)|^2}{\sum_{\xi_i \in \mathcal{I}} |\rho^S_{0,V}(\xi_i)|^2} - \frac{|\rho^S_{0,V}(\psi)|^2 \sum_{\xi_i \in \mathcal{I}} |\rho^S_{1,V}(\xi_i)|^2}{(\sum_{\xi_i \in \mathcal{I}} |\rho^S_{0,V}(\xi_i)|^2)^2} \,, \tag{44}$$

where $\rho^S_{0,V}$ is just the free amplitude. If we take $\mathcal{I}$ to be the one particle sector together with the vacuum state we find

$$P(\psi|\mathcal{I}) = |\rho^S_{1,V}(\psi)|^2 \,. \tag{45}$$

since the free amplitude for the vacuum state is always one independent of the complex structure $J_{\partial M}$ and that for a one particle state vanishes. The first statement can be derived from the expression for the free amplitude of a coherent state in equation (7) and the second from the fact that a one particle state is an odd function and $\rho^S_{n,V}$ is given as a Gaussian integral over an even domain. To get the probability for a whole subset $\mathcal{J} \subseteq \mathcal{I}$ we have to sum that probability again over an orthonormal basis in $\mathcal{J}$.

5.1.1. Unitary evolution

If the boundary of the region M under consideration decomposes as $\partial M = \Sigma_1 \cup \Sigma_2$ and the map T defined in section 3.2.2 exists and is unitary $\hat{\xi} = \xi^R - i\xi^I$ is given explicitly as the complex solution

$$\hat{\xi} = r_2^{-1}\left(\frac{1}{2}(1 - iJ_{\Sigma_2})T\xi_1 + \frac{1}{2}(1 + iJ_{\Sigma_2})\xi_2\right) \tag{46}$$

for all $\xi = (\xi_1, \xi_2) \in L_{\partial M} = L_{\Sigma_1} \times L_{\Sigma_2}$ where r_2 is the homeomorphism defined in section 3.2.2. With equation (10) and using the unitarity of T we can rewrite equation (40)

$$\begin{aligned} \rho^S_\mu(\overline{K}_{\xi_1} \otimes K_{\xi_2}) \quad = \quad & e^{\frac{1}{2}\{\xi_2, T\xi_1\}_{\Sigma_2}} e^{i \int dV(x)\, \mu(x) r_2^{-1}\left(\frac{1}{2}(1-iJ_{\Sigma_2})T\xi_1 + \frac{1}{2}(1+iJ_{\Sigma_2})\xi_2\right)(x)} \times \\ & \times e^{\frac{i}{2}\int dV(x')\, \mu(x')(1-iJ_{\partial M})\eta_D(x')} \,. \end{aligned} \tag{47}$$

Furthermore, we can derive an expression for η_D by using an orthonormal basis of $\mathcal{P}_{\Sigma_2}$ defined in section 3.2.3. We find that $\eta_D = J_{\partial M}(r_1\eta, r_2\eta)$ where $J_{\partial M} = (-T^{-1}J_{\Sigma_2}T, J_{\Sigma_2})$ and

$$\eta(x) := \int d^4x' \mu(x') \sum_{\sigma \in \mathcal{I}} \left(e^-_\sigma(x') e^+_\sigma(x) + e^+_\sigma(x') e^-_\sigma(x)\right) \,. \tag{48}$$

This can be seen by observing that every element $\zeta \in L_M$ can be parameterized as

$$\zeta = \sum_{\sigma \in \mathcal{I}} \left(\zeta_\sigma e^+_\sigma(x) + c.c\right)$$

and it follows that for all $\zeta, \zeta' \in L_M$ holds

$$\begin{aligned} \{\zeta, \zeta'\}_{\Sigma_2} \quad &= \quad i\omega_{\Sigma_2}((1 + iJ_{\Sigma_2})\zeta, (1 - iJ_{\Sigma_2})\zeta) = \sum_{\sigma, \sigma' \in \mathcal{I}} \zeta_\sigma \overline{\zeta'_{\sigma'}} 4i\omega_{\Sigma_2}(e^+_{\sigma'}, e^-_\sigma) \\ &= \quad \sum_{\sigma, \sigma' \in \mathcal{I}} \zeta_\sigma \overline{\zeta'_{\sigma'}} \{e^+_{\sigma'}, e^+_\sigma\}_{\Sigma_2, \mathcal{P}} = \sum_{\sigma \in \mathcal{I}} \zeta_\sigma \overline{\zeta'_\sigma} \,. \end{aligned}$$

Finally, we find that

$$\begin{aligned}2\omega_{\partial M}(\zeta,\eta_D) &= g_{\partial M}(\zeta,\eta) = 2g_{\Sigma_2}(\zeta,\eta)\\ &= \sum_{\sigma\in\mathcal{I}}\left(\zeta_\sigma\overline{\eta_\sigma}+\overline{\zeta_\sigma}\eta_\sigma\right) = \int dV(x)\mu(x)\zeta(x) = D(\zeta)\,. \end{aligned}\tag{49}$$

We obtain

$$\begin{aligned}(1-iJ_{\partial M})\eta_D(x_1,x_2) &= (J_{\partial M}+i)(r_1\eta, r_2\eta)(x_1,x_2)\\ &= 2i\int d^4x'\mu(x')\sum_{\sigma\in\mathcal{I}}(e^-_\sigma(x')e^+_\sigma(x_1), e^+_\sigma(x')e^-_\sigma(x_2))\end{aligned}\tag{50}$$

where x_1 and x_2 are coordinates in neighborhoods of Σ_1 and Σ_2 respectively. η_D is related to the Feynman propagator of the field theory we are investigating [Oec12c].

5.2. The detector model

We now consider coupling a detector on a fixed worldline to the field theory. For first order perturbation theory there is no big difference between modeling the detector as a two level system or a harmonic oscillator as done for example in [LH07]. As the latter will be more convenient for us we will adopt their action principle in what follows. Again to first order this is also similar to a point like atom sitting in an external field.

Having access to this model we can study directly the physical consequences of the choice of complex structure on the transition rates of the detector and use this to study our choices from above in concrete physical situations, as will be done in section 5.4.

The action of [LH07] is given explicitly by

$$S_\gamma(Q) = \int d\tau\frac{m_0}{2}\left((\partial_\tau Q)^2-\Omega_0^2Q^2\right)\,.$$

This is the action for a harmonic oscillator with the frequency Ω_0. It can be seen as a quantum field theory on the worldline $\gamma:\mathbb{R}\to M$ parameterized with proper time τ, i.e. $g_{\mu\nu}(\partial_\tau\gamma)^\mu(\partial_\tau\gamma)^\nu=1$. Holding this in mind we can formulate the corresponding quantum theory using the Schrödinger representation of GBQFT. We find the symplectic potential $[Q,Q']_\tau = m_0Q'\partial_\tau Q|_\tau$ and the symplectic form $\omega_\tau(Q,Q') = \frac{m_0}{2}\left(Q'\partial_\tau Q - Q\partial_\tau Q'\right)|_\tau$. For the complex structure we choose

$$J_\tau := \frac{\partial_\tau}{\sqrt{-\partial_\tau^2}}\,.$$

We consider a region $\gamma_0 := [\tau_1,\tau_2]$ that is a subset of the worldline γ. The solutions of the equations of motion that form the solution space L_{γ_0} can be parameterized as

$$Q(\tau) = \frac{1}{\sqrt{2m_0\Omega_0}}\left(qe^{-i\Omega_0\tau}+\overline{q}e^{i\Omega_0\tau}\right)\,.$$

With $J_{\partial\gamma_0} = (-J_\tau, J_\tau)$ the complex structure on $L_{\partial\gamma_0} = L_{\overline{\tau_1}}\otimes L_{\tau_2}$ we find using equation (50) with $e^+ = (2m_0\Omega_0)^{-1/2}e^{-i\Omega_0\tau}$ and $e^- = (2m_0\Omega_0)^{-1/2}e^{i\Omega_0\tau}$ that

$$(1-iJ_{\partial\gamma_0})\eta_D(\tau) = \int d\tau'\,\mu(\tau')G_F(\tau',\tau)\,.\tag{51}$$

which is a solution of the homogeneous equation near the boundary and thus determines two solutions of the homogeneous equations that are joined by a solution to the inhomogeneous equation in the interior.

$$G_F(\tau',\tau) := \frac{i}{m_0\Omega_0}\left(\theta(\tau-\tau')e^{-i\Omega_0(\tau'-\tau)}+\theta(\tau'-\tau)e^{i\Omega_0(\tau'-\tau)}\right)$$

is the Feynman propagator for our one dimensional field theory. For the amplitude we obtain for the boundary state $K_{(\Theta_1,\Theta_2)} = \overline{K}_{\Theta_1} \otimes K_{\Theta_2}$ with

$$\Theta_i(\tau) = \frac{1}{\sqrt{2m_0\Omega_0}} \left(\vartheta_i e^{-i\Omega_0\tau} + \overline{\vartheta_i} e^{i\Omega_0\tau} \right) .$$

from (47) the expression

$$\begin{aligned} & \rho^S_{\mu_Q}(\overline{K}_{\Theta_1} \otimes K_{T\Theta_2}) \\ = \; & \exp\left(\frac{1}{2}\{\Theta_1, \Theta_2\}_{\tau_2} + i \int d\tau \, \mu(\tau) \left(\frac{1}{2}(1 - iJ_\tau)\Theta_1(\tau) + \frac{1}{2}(1 + iJ_\tau)\Theta_2(\tau) \right) \right. \\ & \left. + \frac{i}{2} \int d\tau d\tau' \, \mu(\tau)\mu(\tau') G_F(\tau, \tau') \right) \\ = \; & \exp\left(\frac{1}{2}\vartheta_1\overline{\vartheta_2} + \frac{i}{\sqrt{2m_0\Omega_0}} \int d\tau \, \mu(\tau) \left(\overline{\vartheta_1} e^{i\Omega_0\tau} + \vartheta_2 e^{-i\Omega_0\tau} \right) \right. \\ & \left. + \frac{i}{2} \int d\tau d\tau' \, \mu(\tau)\mu(\tau') G_F(\tau, \tau') \right) \end{aligned}$$

To turn the quantized harmonic oscillator into a detector we still need an interaction term. For the interaction with a scalar field in four dimensions it is given as [LH07]

$$\mathcal{V}_{UD}(\phi, Q) = \lambda \int dV(x) \int d\tau \, \rho^{-1} \delta^{(4)}(x - \gamma(\tau)) \phi(x) Q(\tau) . \tag{52}$$

where ρ is defined by $dV(x) = \omega_{\mu_1 \cdots \mu_4} dx^{\mu_1} \cdots dx^{\mu_3} = \rho \, \epsilon_{\mu_0 \cdots \mu_3} dx^{\mu_1} \cdots dx^{\mu_3}$ with ϵ the totally antisymmetric Levi-Civita tensor. We want to investigate the interacting theory (52) in a region in Minkowski space with flat timelike or spacelike boundaries. We assume that the map T exists and that it is unitary. In first order perturbation theory we find for the coherent states K_ξ and $K_\Theta = \overline{K}_{\Theta_1} \otimes K_{\Theta_2}$ for the field and the detector respectively that

$$\begin{aligned} & \rho^S_{1,V}(K_\xi \otimes K_\Theta) \\ := \; & \int dV(x) \int d\tau \, \mathcal{V}_{UD}\left(-i\frac{\delta}{\delta\mu_\phi(x)}, -i\frac{\delta}{\delta\mu_Q(\tau)} \right) \rho^S_{\mu_\phi}(K_\xi) \rho^S_{\mu_Q}(K_\Theta) \Big|_{\mu_\phi, \mu_Q = 0} , \end{aligned}$$

where $\rho^S_{\mu_\phi}(K_\xi)$ and $\rho^S_{\mu_Q}(K_\Theta)$ are the free amplitudes of the interacting quantum systems. With $\rho^S_{\mu_\phi}(K_\xi)$ by equation (40) given as

$$\rho^S_{\mu_\phi}(K_\xi) = \rho_M(K_\xi) e^{iD(\hat{\xi}) + \frac{i}{2} D((1 - iJ_{\partial M})\eta_D)} \tag{53}$$

and the fact that the only term of first order in μ_ϕ is $D(\hat{\xi})$ we obtain with the definition of the potential in equation (52)

$$\begin{aligned} & \rho^S_{1,V}(K_\xi \otimes K_\Theta) \\ = \; & -\lambda \rho_{\gamma_0}(K_\Theta) \rho_M(K_\xi) \int d\tau \, \hat{\Theta}(\tau) \hat{\xi}(\gamma(\tau)) \\ = \; & -\lambda \rho^S(K_\xi) e^{\frac{1}{2}\vartheta_1\overline{\vartheta_2}} \frac{1}{\sqrt{2m_0\Omega_0}} \int d\tau \, \left(\overline{\vartheta_1} e^{i\Omega_0\tau} + \vartheta_2 e^{-i\Omega_0\tau} \right) \hat{\xi}(\gamma(\tau)) . \end{aligned}$$

For a one particle state, we derive

$$\begin{aligned} & \rho^S_{1,V}(a^\dagger_\xi K_0 \otimes a^\dagger_\Theta K_0) = 2\frac{d}{d\alpha}\frac{d}{d\beta} \rho^S_{1,V}(K_{\alpha\xi} \otimes K_{\beta\Theta}) \Big|_{\alpha,\beta=0} \\ = \; & \frac{\sqrt{2}\lambda}{\sqrt{m_0\Omega_0}} \int_{\tau_1}^{\tau_2} d\tau \, \left(\overline{\vartheta_1} e^{i\Omega_0\tau} + \vartheta_2 e^{-i\Omega_0\tau} \right) \hat{\xi}(\gamma(\tau)) . \end{aligned}$$

From equation (45) we know that the absolute square of $\rho^S_{1,V}(a^\dagger_\xi K_0 \otimes a^\dagger_\Theta K_0)$ can then be interpreted as a transition probability in the one particle sector. The transitions that are interesting for us are the transition of the detector from the ground state to an excited state and vice versa. So we set $\vartheta_1 = 0$ and $\vartheta_2 = 1$ and find

$$P^{\Omega_0,\xi}(g \to e) \quad := \quad \frac{2\lambda^2}{m_0\Omega_0}\int_{\tau_1}^{\tau_2} d\tau d\tau' \, e^{-i\Omega_0(\tau-\tau')}\hat{\xi}(\gamma(\tau))\overline{\hat{\xi}(\gamma(\tau'))}\,, \tag{54}$$

and with $\vartheta_1 = 1$ and $\vartheta_2 = 0$ we obtain

$$P^{\Omega_0,\xi}(e \to g) \quad := \quad \frac{2\lambda^2}{m_0\Omega_0}\int_{\tau_1}^{\tau_2} d\tau d\tau' \, e^{i\Omega_0(\tau-\tau')}\hat{\xi}(\gamma(\tau))\overline{\hat{\xi}(\gamma(\tau'))}\,. \tag{55}$$

The map $\xi \mapsto \hat{\xi} = \xi^R - i\xi^I$ is a linear map. Hence, for every two orthonormal elements $\xi, \xi' \in L_{\partial M}$ the sum $P_\xi(g \to e) + P_{\xi'}(g \to e)$ is invariant under a transformation to the orthonormal states labeled by $a\xi + b\xi'$ and $b\xi - a\xi'$ where $a, b \in \mathbb{R}$, i.e.

$$P^{\Omega_0,\xi}(g \to e) + P^{\Omega_0,\xi'}(g \to e) = P^{\Omega_0,a\xi+b\xi'}(g \to e) + P^{\Omega_0,b\xi-a\xi'}(g \to e)\,. \tag{56}$$

where $a^2 + b^2 = 1$. We find that summing the probability in equation (54) or (55) over a general subset of a basis in $L_{\partial M}$ we arrive at an expression invariant under a general orthogonal basis transformation. This leads to the generalization of Fermi's golden rule to general regions. We sum over an orthonormal basis in the set of states that represent the knowledge we have about the system [Oec08].

5.3. The Unruh-DeWitt detector no-click condition and the energy-momentum flux for timelike hyperplanes

In this section we show how the detector response can be used to formulate a condition that determines the complex structure for propagating waves when claiming the complex structure to possess all the symmetries of the spacetime region. We are dealing with $1+1$ dimensional Minkowski space and a region $M = \Sigma \times [x_l, x_r]$ bounded by parallel timelike hyperplanes Σ_l and Σ_r where Σ_r is oriented as Σ and Σ_l has opposite orientation. Hence, the symmetries in the global coordinates in which the time direction is tangential to the timelike hyperplanes and the metric takes the form $g = \mathrm{diag}(1, -1)$ at every point of M are: time translation and time reversal. We use exactly this coordinate system in the following. We assume the Unruh-DeWitt detector to be at rest, i.e. $\gamma(\tau) = (\tau, z)$ inside the region M throughout this whole section (see figure 5.3).

The additional ingredient for determining the complex structure is the positive flux condition formulated in section 4.3.2. There we already introduced the terms of positive flux modes as elements of the space $L^{\mathbb{C}}_{\partial M,+} \subset L^{\mathbb{C}}_{\partial M}$ spanned by energy eigenstates ξ_{p_0} fulfilling

$$n_\mu T^{\mu 0}(\mathfrak{Re}(\xi_{p_0}))(x) > 0 \tag{57}$$

at every point $x \in \partial M$ with n the unit normal covector field to ∂M pointing out of the region M. Now, to determine the complex structure we require that the set of states corresponding to positive flux modes should give a vanishing transition rate of the detector. To make this mathematically clear we first assume that we are dealing with a unitary complex structure. Since the complex structure respects the symmetries of the classical configuration we can deal with all the energy subspaces of $L^{\mathbb{C}}_\Sigma$ separately. To determine the complex structure we look at the space $L^{\mathbb{C},E}_\Sigma := L^{\mathbb{C},p_0}_\Sigma \oplus L^{\mathbb{C},-p_0}_\Sigma$ with $L^{\mathbb{C},p_0}_\Sigma = \{\sum_{\sigma_1=\pm} a_{\sigma_1} e(p_0, \sigma_1 p_1) | \, a_{\sigma_1} \in \mathbb{C}\}$ where $e(p_0, p_1) = e^{-i(p_0 x_0 - p_1 x_1)}$. Using this split we investigate the no-click condition for elements of $L^{\mathbb{C}}_{\partial M,+}$ of the form $\xi^{\mathbb{C}}_+ = (\xi_l, \xi_r)$ with $\xi_l = c_{l,+}e(p_0, -p_1) + c_{l,-}e(-p_0, p_1)$ and $\xi_r = c_{r,+}e(p_0, p_1) + c_{r,-}e(-p_0, -p_1)$. We define the boundary configuration $\xi_+ := \mathfrak{Re}(\xi^{\mathbb{C}}_+) \in L_{\partial M}$. Let the coefficients $c_{i,\pm}$ be given such that ξ_+ corresponds to a



Figure 2: Schematic picture of the region M containing the worldline γ of the detector at rest. The horizontal axis represents space and the vertical axis time.

normalized state $a^\dagger_{\xi_+} K_0$, we find that for the field being in this state the probability for the transition of a detector at rest, i.e. worldline $\gamma(\tau) = (\tau, z)$ from the ground state to the excited state following from equation (54) is given as

$$P_\eta(g \to e) \quad = \quad \frac{2\lambda^2}{m_0 \Omega_0} \int_{-\infty}^{\infty} d\tau d\tau' \, e^{-i\Omega_0(\tau - \tau')} \hat{\xi}_+(\tau, z) \overline{\hat{\xi}_+(\tau', z)} \,, \tag{58}$$

where

$$\hat{\xi}_+ \quad = \quad \xi^R_+ - i\xi^I_+ = \frac{1}{2}\left(1 - iJ_{\Sigma_r}\right) \mathfrak{Re}(\xi_l) + \frac{1}{2}\left(1 + iJ_{\Sigma_r}\right) \mathfrak{Re}(\xi_r) \,.$$

The no-click condition is required to hold for every element of $L^{\mathbb{C}}_{\partial M,+}$, that is for every set of coefficients $c_{l,\pm}$ and $c_{r,\pm}$. Considering the special case of $c_{r,+} = c_{r,-}$ and $c_{l,+} = c_{l,-} = 0$ we obtain that

$$\hat{\xi}_+ \quad = \quad \xi^R_+ - i\xi^I_+ = c_{r,+} \left(\frac{1}{2}\left(1 + iJ_{\Sigma_r}\right) e(p_0, p_1) + \frac{1}{2}\left(1 + iJ_{\Sigma_r}\right) e(-p_0, -p_1) \right) .$$

which means that for the detector not to click we need $(1 + iJ_{\Sigma_r})\, e(-p_0, -p_1) = 0$ which leads to $J_{\Sigma_r} e(-p_0, -p_1) = ie(-p_0, -p_1)$ and with complex conjugation $J_{\Sigma_r} \overline{e(-p_0, -p_1)} = J_{\Sigma_r} e(p_0, p_1) = -ie(p_0, p_1)$. Together with the time reversal invariance which tells us that $J_{\Sigma_r} e(p_0, -p_1) = ie(p_0, -p_1)$ and $J_{\Sigma_r} e(-p_0, p_1) = -ie(-p_0, p_1)$ this already determines the complex structure completely. The one that is determined is the one we obtained in 4.3.1. We obtain that

$$\hat{\xi}_+ = c_{l,+} e(p_0, p_1) + c_{r,+} e(p_0, -p_1) \propto e^{-ip_0 x_0} \,,$$

such that the detector does not click for every choice of $c_{l,\pm}$ and $c_{r,\pm}$.

5.4. An explicit experimental situation

In the preceding sections we saw that also on timelike hypersurfaces reasonable conditions for the complex structure can be imposed. For propagating waves all of them lead to a unique complex structure that, at least for hyperplanes coincide for the different conditions. For evanescent waves instead we found that no complex structure exists that fulfills the corresponding conditions.

In this section we use a different approach to deal with evanescent waves. We show that the complex structure can and must be determined depending on the specific physical situation outside of the quantization region M when evanescent waves can be prepared. In particular, we canonically quantize a system in which evanescent waves appear on spacelike hypersurfaces and use the results to determine the complex structure for timelike hypersurfaces.

5.4.1. Canonical quantization in the presence of dielectric slabs

An example of a physical situation in which evanescent waves can be prepared is Maxwell electrodynamics in the presence of a dielectric slab. This was investigated in [CM71] where the authors model the experimental situation as two half spaces of different refractive index with one of them being the vacuum refractive index. In this setup solutions exist that are propagating waves in the dielectric medium and become evanescent in the vacuum. The authors of [CM71] find after quantization that their theoretical results coincide with semiclassical calculations and in a later publication [CMD72] they also show that they coincide with experiments. We are going to consider a similar setup for the massive scalar field in $1+1$ dimensions. This toy model recovers the basic situation found in [CM71] but makes it much easier to understand the calculations which will allow us to focus on the basic insights concerning evanescent waves. We will use the Unruh-DeWitt detector as a toy model for the atoms in the electromagnetic field near the dielectric boundary. Throughout this section we will again assume the detector to be at rest, i.e. $\gamma(\tau) = (\tau, z)$.



Figure 3: Schematic picture for the preparation of evanescent waves in a toy model of $1+1$-dimensional massive scalar field theory. The horizontal axis represents space and the vertical axis time.

The setup we consider is shown in figure 5.4.1. The $1+1$ dimensional spacetime is separated in three regions with different metric tensor g. For region I and III we have

$$g = \begin{pmatrix} \frac{1}{n} & 0 \\ 0 & -1 \end{pmatrix}$$

and for region II we have:

$$g = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

The boundaries of the regions are at ξ and ξ'. The conditions for the construction of solutions ϕ of the Klein-Gordon equation

$$\left(\partial_\mu \sqrt{g}\, g^{\mu\nu} \partial_\nu + \sqrt{g}\, m^2\right) \phi(x) = 0\,, \tag{59}$$

at ξ and ξ' are the continuity of ϕ and its first derivative. With the separation

$$\phi^{\pm}_{p_0}(x) = \varphi^{\pm}_{p_0}(x_1) e^{i p_0 x_0} \tag{60}$$

we obtain for $p_0^2 < m^2$ the set of solutions

$$\varphi^+_{p_0}(x_1) = \begin{cases} N_{p_0} e^{-ipx_1} & \text{for} \quad x_1 < -\xi \\ N_{p_0} e^{ip\xi} \Big(\cosh p_1(x_1+\xi) - i\frac{p}{p_1} \sinh p_1(x_1+\xi)\Big) & \text{for} \quad -\xi \leq x_1 \leq \xi \\ e^{-ipx_1} + C_{p_0} e^{ipx_1} & \text{for} \quad \xi < x_1 \end{cases} , \tag{61}$$

and $\varphi^-_{p_0}(x_1) = \varphi^+_{p_0}(-x_1)$. The constants are given as

$$\begin{aligned} N_{p_0} &= e^{-2ip\xi} \left(\cosh 2p_1\xi - \frac{i}{2}\left(\frac{p}{p_1} - \frac{p_1}{p}\right) \sinh 2p_1\xi\right)^{-1} \\ C_{p_0} &= -N_{p_0} \frac{i}{2}\left(\frac{p}{p_1} + \frac{p_1}{p}\right) \sinh 2p_1\xi \,, \end{aligned} \tag{62}$$

where $p = (np_0^2 - m^2)^{1/2}$ and $p_1 = (m^2 - p_0^2)^{1/2}$. These solutions are chosen such that with the symplectic form for an equal time hypersurface given as

$$\omega_{x_0}(\phi, \phi') = \frac{1}{2}\int_{-\infty}^{\infty} dx_1 \, \sqrt{g}\, g^{00} (\phi\partial_0\phi' - \phi'\partial_0\phi)\big|_{x_0} \,.$$

they fulfill the orthonormality relation with the Klein-Gordon inner product

$$(\phi^s_{p_0}, \phi^{s'}_{p'_0})_{KG} := i\omega_M(\overline{\phi^s_{p_0}}, \phi^{s'}_{p'_0}) = 2\pi\sqrt{n} p_0 \delta(p - p')\delta_{s,s'} \,. \tag{63}$$

This is shown in appendix A.3. With equation (63) the canonical quantization prescription leads to the absorption and emission operator

$$\hat{\phi}(x) = \int_0^{\infty} \frac{dp}{2\pi\sqrt{n}p_0} \sum_{s=\pm} \big(a_{p,s}\phi^s_{p_0}(x) + h.c.\big)$$

with the commutation relations $[a^\dagger_{p,s}, a_{p',s'}] = (\phi^s_{p_0}, \phi^{s'}_{p'_0})_{KG} = 2\pi\sqrt{n}p_0\delta(p - p')\delta_{s,s'}$. The total probabilities for an Unruh-DeWitt detector with the world line $\gamma(\tau) = (\tau, z)$ to emit a field particle while passing from its ground state to an exited state and vice versa is then given as

$$\begin{aligned} P^{\Omega_0}(g \to e) &= \frac{\lambda^2}{m_0\Omega_0} \int \frac{dp}{2\pi\sqrt{n}p_0} \int_{-\infty}^{\infty} d\tau d\tau' e^{-i(\Omega_0 + p_0)(\tau - \tau')} \sum_{s=\pm} |\varphi^s_{p_0}(z)|^2 \\ P^{\Omega_0}(e \to g) &= \frac{\lambda^2}{m_0\Omega_0} \int \frac{dp}{2\pi\sqrt{n}p_0} \int_{-\infty}^{\infty} d\tau d\tau' e^{i(\Omega_0 - p_0)(\tau - \tau')} \sum_{s=\pm} |\varphi^s_{p_0}(z)|^2 \,. \end{aligned} \tag{64}$$

It turns out that the probability $P^{\Omega_0}(g \to e)$ vanishes and with the expression in equation (61) we find the transition probability

$$\begin{aligned} P^{\Omega_0}(e \to g) \quad &:= \quad \frac{\lambda^2\sqrt{n}}{m_0\Omega_0 p} \sum_{s=\pm} |\varphi^s_{\Omega_0}(z)|^2 \\ &= \quad \frac{\lambda^2\sqrt{n}}{m_0\Omega_0 p} |N_{\Omega_0}|^2 \Bigg(\left(\frac{p_1}{p} - \frac{p}{p_1}\right) \\ &\qquad + \frac{1}{2}\left(\frac{p_1}{p} + \frac{p}{p_1}\right) \big(\cosh 2p_1(z + \xi) + \cosh 2p_1(z - \xi)\big)\Bigg) \delta(0)^2 \\ &= \quad \frac{\lambda^2\sqrt{n}}{m_0\Omega_0 p} |N_{\Omega_0}|^2 \left(\left(\frac{p_1}{p} - \frac{p}{p_1}\right) + \left(\frac{p_1}{p} + \frac{p}{p_1}\right) \cosh 2p_1 z \, \cosh 2p_1\xi\right) , \end{aligned} \tag{65}$$

that depends on the space point z at which the detector rests and on the energy Ω_0. If we want to know the probabilities for the absorption of a particle by the detector undergoing the same transitions as above we just have to interchange Ω_0 by $-\Omega_0$ in equation (64). Then we find that $P^{\Omega_0}_{\text{absorption}}(e \to g) = 0$ and $P^{\Omega_0}_{\text{absorption}}(g \to e) = P^{\Omega_0}_{\text{emission}}(e \to g)$ where $P^{\Omega_0}_{\text{emission}}(e \to g)$ is the transition probability in equation (64).

We will now repeat this calculation in GBQFT, by making use of the tools developed in the foregoing sections. We choose the region $M = \Sigma \times [x_l, x_r]$ to lie completely in the interior region II which means that the classical field theory inside the region M and in a neighborhood of ∂M is just that of $1+1$-dimensional Minkowski space. For M being extended to both timelike infinities and for an Unruh-DeWitt detector at rest inside the region M we want to recover all the first order results of the canonical quantization. Of course, we have to stress what we mean by recovering the results of the canonical theory. We will explain this in the following.

5.4.2. The canonical case in GBQFT

First, we make contact with the GBQFT considering a configuration with spacelike hypersurfaces: Already in section 4.3.2 we used that for spacelike hypersurfaces and unitary evolution we recover standard geometrical quantized field theory which can be seen from the expressions in section 3.2.2. Let us consider the region $M' = \Sigma \times [t_i, t_f]$ with the boundaries Σ_i and Σ_f with the orientations pointing outside the region M'. The split of the boundary Hilbert space as $\mathcal{H}_{\partial M'} = \mathcal{H}_{\Sigma_i} \otimes \mathcal{H}_{\Sigma_f}$ and the existence of a the unitary map $U : \mathcal{H}_{\Sigma_i} \to \mathcal{H}_{\Sigma_f}$ with $U(\psi)(\phi) = \psi(T^{-1}\phi)$ for all $\psi \in \mathcal{H}_{\Sigma_i}$ leads to an identification of the Hilbert spaces $\mathcal{H}_{\Sigma_i}$ and $\mathcal{H}_{\Sigma_f}$ and the reduction of the amplitude map to the inner product. It is then easy to see that the canonical theory is completely equivalent to GBQFT with the complex structure

$$J_{\Sigma_f} = \frac{\partial_0}{\sqrt{-\partial_0^2}} \tag{66}$$

as can be found in section 4.1.1. As we already argued in section 4.3.2 in the GBQFT we find a slightly different picture of the states on the boundary. The states on the future boundary and those on the past boundary have no a priori roles in the theory and they have to be interpreted. For that purpose we use the Unruh-DeWitt detector. Let t_i and t_f go to $-\infty$ and ∞ respectively. We obtain for an arbitrary boundary one particle state $a^\dagger_\xi K_0 \in \mathcal{H}_{1,\partial M'} \cong L_{\partial M'}$ with $\xi = (\xi_i, \xi_f) \in L_{\partial M}$ the transition probability

$$P^{\Omega_0,\xi}(g \to e) = \frac{2\lambda^2}{m_0 \Omega_0 p_1} \int_{-\infty}^{\infty} d\tau d\tau'\, e^{-i\Omega_0(\tau-\tau')} \hat{\xi}(\gamma(\tau)) \overline{\hat{\xi}(\gamma(\tau'))}\,. \tag{67}$$

where $\hat{\xi} = \frac{1}{2}(1 - iJ_{\Sigma_f})\xi_i + \frac{1}{2}(1 + iJ_{\Sigma_f})\xi_f$. Investigating this in the wave function basis $\{\phi^\pm_{p_0}\}$ defined in (60) which consists of plane waves in region I and III we find that the one particle Hilbert space $\mathcal{H}_{1,\partial M'}$ splits into an orthogonal direct sum $\mathcal{H}_{1,\partial M'} = \mathcal{H}^{\text{out}}_{1,\partial M'} \oplus \mathcal{H}^{\text{in}}_{1,\partial M'}$ where $\mathcal{H}^{\text{out}}_{1,\partial M'}$ is the set of all those one particle states corresponding to a boundary solution $\xi \in L_{\partial M'}$ with $\hat{\xi}$ an element of the space spanned by the basis $\{\phi^\pm_{p_0}\}$. $\mathcal{H}^{\text{in}}_{1,\partial M'}$ is defined analogously as the set of states corresponding to boundary solutions with $\hat{\xi}$ in $\text{span}\{\overline{\phi^\pm_{p_0}}\}$. This can be seen the following way: The map $\hat{}: L_{\partial M'} \to L^{\mathbb{C}}_{\tilde{M}'}$, $\xi = \xi^R + J_{\partial M'}\xi^I \mapsto \hat{\xi}$ can be inverted as $\hat{}^{-1} : L^{\mathbb{C}}_{\tilde{M}'} \to L_{\partial M}$, $\mathfrak{Re}\,\eta + i\mathfrak{Im}\,\eta \mapsto \mathfrak{Re}\,\eta - J_{\partial M'}\mathfrak{Im}\,\eta$. Using this insight we can write the inner product of two one particle states labeled by ξ and ξ' as

$$\langle a^\dagger_\xi K_0, a^\dagger_{\xi'} K_0 \rangle = \{\xi', \xi\}_{\partial M'} = g_{\partial M'}(\hat{\xi}', \overline{\hat{\xi}})\,. \tag{68}$$

Obviously states corresponding to $\text{span}\{\overline{\phi^\pm_{p_0}}\}$ are orthogonal to those in $\text{span}\{\phi^\pm_{p_0}\}$.

We find that summing the transition probability (67) and the reverse process over an orthonormal basis in $\mathcal{H}^{\text{out}}_{1,\partial M'}$ we obtain a vanishing probability $P^{\Omega_0,\text{out}}(g \to e)$ and $P^{\Omega_0,\text{out}}(e \to g)$ as in equation (64) respectively. For an orthonormal basis in $\mathcal{H}^{\text{in}}_{1,\partial M'}$ we find that $P^{\Omega_0,\text{in}}(e \to g)$ vanishes and

$P^{\Omega_0,\mathrm{in}}(g \to e)$ recovers the expression in equation (64). From the definition of the map $\hat{}$ we obtain that the states in $\mathcal{H}^{\mathrm{out}}_{1,\partial M'}$ are exactly those corresponding to boundary configurations that are zero on the initial hypersurface Σ_i and nonzero on Σ_f and for $\mathcal{H}^{\mathrm{in}}_{1,\partial M'}$ vice versa. Let us recall here again what we call the canonical result: The one particle Hilbert space can be split in an orthonormal sum of outgoing and ingoing states where when summed over an orthonormal basis of the outgoing states we find that for every Ω_0 the probability $P^{\Omega_0,\mathrm{out}}(g \to e)$ vanishes and that $P^{\Omega_0,\mathrm{out}}(e \to g)$ is given by (64) and for the in states vice versa. This is what we want to recover for timelike hypersurfaces, for every transition energy $\Omega_0 > 0$.

5.4.3. Timelike hyperplanes in GBQFT

Let us now come back to the region $M = \Sigma \times [x_l, x_r]$ with timelike boundaries Σ_l and Σ_r with unit normal pointing to the outside of M and M lying completely in region II. The symplectic form is given as $\omega_{\partial M} = \omega_{\Sigma_l} + \omega_{\Sigma_r}$ where

$$\omega_{\Sigma_{l,r}}(\eta_1, \eta_2) = \sigma_{l,r} \frac{1}{2} \int \mathrm{d}x_0 \, \mathrm{d}^2\tilde{x} \ \left(\eta_1(t, x_1, \tilde{x}) \partial_1 \eta_2(t, x_1, \tilde{x}) - \eta_2(t, x_1, \tilde{x}) \partial_1 \eta_1(t, x_1, \tilde{x})\right)|_{\Sigma_{l,r}} . \quad (69)$$

and $\sigma_l = -1$ and $\sigma_r = 1$. Since the boundary of M is a disjoint union $\partial M = \Sigma_l \cup \Sigma_r$ the complex structure $J_{\partial M}$ acts as complex structures J_{Σ_l} and J_{Σ_r} on L_{Σ_l} and L_{Σ_r} respectively.

For this region we want to recover the canonical result. First of all the sum $P^{\Omega_0,\mathrm{tot}}(g \to e) := P^{\Omega_0,\mathrm{out}}(g \to e) + P^{\Omega_0,\mathrm{in}}(g \to e)$ is a sum over an orthonormal basis in the one particle Hilbert space which is invariant under a change of basis as we explained at the end of section 5.2. But then $P^{\Omega_0,\mathrm{tot}}(g \to e)$ cannot be modified by a suitable choice of boundary states, on the other hand $P^{\Omega_0,\mathrm{tot}}(g \to e)$ is still equal to (65) which depends on the distance ξ to regions I and III and on the refraction index n, in other words on information about the physics outside of M. The only way to recover the canonical result is by selecting an appropriate complex structure which encodes this physical information. This is what we shall do now.

More precisely, in the following we show that for the infinitely extended region M bounded by timelike hyperplanes lying completely in the region II as it can be seen in figure 4 the complex structure can be determined completely by the condition to recover the full canonical result when we assume that it is unitary and possesses the symmetries of region M, that is time translation and time reversal invariance.



Figure 4: The region M lies completely inside the region II and the detector is at rest inside the region M. The horizontal axis represents space and the vertical axis time.

In fact, in the situation considered here where the complex structure is required to be invariant under time translation it suffices to investigate the transition probability just for the outgoing states: Suppose we found a subspace $\mathcal{H}^{\text{out}}_{1,\partial M} \subset \mathcal{H}_{1,\partial M}$ such that

$$\sum_{\xi \in \mathcal{H}^{\text{out}}_{1,\partial M}} P^{\Omega_0,\xi}(g \to e) = 0,$$

where the sum is over an ONB of $\mathcal{H}^{\text{out}}_{\partial M}$, and the same equation with $(g \to e)$ replaced by $(e \to g)$ yields (64), in other words the outgoing states recover their canonical result. (67) implies that $P^{\Omega_0,\xi}(g \to e) = P^{\Omega_0,\bar{\xi}}(e \to g)$ where $\bar{\xi}$ is the real boundary configuration corresponding to $\overline{\hat{\xi}}$. Hence, if we define $\mathcal{H}^{\text{in}}_{1,\partial M}$ as the set of all elements $a^\dagger_{\bar{\xi}} K_0 \in \mathcal{H}_{\partial M}$ such that $a^\dagger_{\xi} K_0 \in \mathcal{H}^{\text{out}}_{1,\partial M}$, then the ingoing states recover their canonical result. We identify the set of outgoing states as those that correspond to complex solutions $\hat{\xi} \propto e^{-ip_0 x_0}$ with $p_0 \geq 0$, and then the ingoing states are the ones with $p_0 < 0$. Every element of $\mathcal{H}_{1,\partial M}$ can be written as a linear combination of those states, hence $\mathcal{H}_{1,\partial M} = \mathcal{H}^{\text{out}}_{1,\partial M} \cup \mathcal{H}^{\text{in}}_{1,\partial M}$. Moreover, one can check that $\mathcal{H}^{\text{out}}_{1,\partial M}$ and $\mathcal{H}^{\text{in}}_{1,\partial M}$ are isotropic subspaces with respect to $\omega_{\partial M}$, and therefore $\mathcal{H}^{\text{out}}_{1,\partial M}$ and $\mathcal{H}^{\text{in}}_{1,\partial M}$ are orthogonal iff $J_{\partial M}$ leaves both subspaces invariant: For $\xi, \xi' \in \mathcal{H}^{\text{out}}_{1,\partial M}$,

$$\langle a^\dagger_{\xi} K_0, a^\dagger_{\bar{\xi}} K_0 \rangle = g_{\partial M}\left(\hat{\xi}, \hat{\xi}'\right) = \omega_{\partial M}\left(\hat{\xi}, J_{\partial M}\hat{\xi}'\right)$$

is zero iff $J_{\partial M}\xi'$ is again in $\mathcal{H}^{\text{out}}_{1,\partial M}$. But this is of course the case if $J_{\partial M}$ is invariant under time translation, for then it must commute with the time translation operator $E := i\partial_0$. Then $J_{\partial M}$ leaves the eigenspaces of E invariant.

There are even more conclusions we can draw from the time translation invariance: If $J_{\partial M}$ leaves the eigenspaces of E invariant we can talk about a complex structure for every energy subspace $L^{p_0}_{\partial M} := \{\mathfrak{Re}(c_l e^+_{p_0;\Sigma_l}, c_r e^+_{p_0;\Sigma_r}) | \, c_l, c_r \in \mathbb{C}\}$ separately. We do this by writing the eigenfunctions of J_{Σ_r} in the form

$$e^+_{p_0;\Sigma_r} := (c_a(p_0)\cosh(p_1 x_1) + c_b(p_0)\sinh(p_1 x_1))e^{-ip_0 x_0}\,, \tag{70}$$

with $p_1 = \sqrt{m^2 - p_0^2}$ and eigenfunctions J_{Σ_l} analogously. The choice of complex structure per energy subspace is now reduced to the choice of coefficients $c_a(p_0), c_b(p_0) \in \mathbb{C}$. Of course, the definition of a complex structure by (70) stays invariant if we multiply the expression in (70) with an additional energy dependent complex factor. Hence, the only thing that matters is the quotient $c_a(p_0)/c_b(p_0)$.

Using this result, we obtain that two states corresponding to $\xi_{p_0} \in L^{p_0}_{\partial M}$ and $\xi_{p'_0} \in L^{p'_0}_{\partial M}$ with $p_0 \neq p'_0$ are orthogonal. This can be seen in the following way: Take two states of the form $\xi_{p_0} = \mathfrak{Re}(c_l(p_0)e^+_{p_0;\Sigma_l}, c_r(p_0)e^+_{p_0;\Sigma_r})$. We find that

$$\begin{aligned}
\{\xi_{p_0}, \xi_{p'_0}\}_{\partial M} &= i\omega_{\partial M}((1 + iJ_{\partial M})\xi_{p_0}, (1 - iJ_{\partial M})\xi_{p'_0}) \\
&= i\omega_{\Sigma_l}(c_l(p_0)e^+_{p_0;\Sigma_l}, \overline{c_l(p'_0)e^+_{p'_0;\Sigma_l}}) + i\omega_{\Sigma_r}(c_r(p_0)e^+_{p_0;\Sigma_r}, \overline{c_r(p'_0)e^+_{p'_0;\Sigma_r}}) \propto \delta(p_0 - p'_0)\,.
\end{aligned}$$

Hence, $\{\xi_{p_0}, \xi_{p'_0}\}_{\partial M} = 0$ for all elements in $\xi_{p_0} \in L^{p_0}_{\partial M}$ and $\xi_{p'_0} \in L^{p'_0}_{\partial M}$ with $p_0 \neq p'_0$.

Now, we want to recover the canonical result: We see from the transition probability in (54) and (55) that the detector with transition energy Ω_0 is only sensible to states in $\mathcal{H}^{\Omega_0}_{1,\partial M} \otimes \mathcal{H}^{-\Omega_0}_{1,\partial M}$. We can use the orthogonality of different energy subspaces and the reduction of the decomposition problem to the outgoing states due to the the time translation invariance. That reduces the task of finding a decomposition of the one particle Hilbert space in out- and ingoing states to the task of finding two orthogonal states for every $E = |p_0|$ spanning the orthogonal subspace $\mathcal{H}^{\text{out},E}_{1,\partial M}$, that is the set of states $a^\dagger_{\xi} K_0 \in \mathcal{H}^{|p_0|}_1 := \mathcal{H}^{p_0}_{1,\partial M} \otimes \mathcal{H}^{-p_0}_{1,\partial M}$ such that $\hat{\xi} \propto e^{-ip_0 x_0}$ with $p_0 > 0$.

From equation (68) we know that this is equivalent to the problem of finding two solutions ξ_1, ξ_2 for every p_0 in the complexification $L^{p_0,\mathbb{C}}_{\tilde{M}}$ such that they are orthonormal with respect to the sesquilinear map $(\cdot,\cdot) := g_{\partial M}(\cdot,\bar{\cdot})$ which is antilinear in the second entry. We start with the two linearly independent eigenfunctions of J_{Σ_r} (see 70) defining $\xi^+_{p_0} := e^+_{p_0;\Sigma_r}$ and $\xi^-_{p_0} := \overline{\xi^+_{-p_0}}$. We define the normalization constant $c^+_{p_0}$ such that $(\xi^+_{p_0}, \xi^+_{p_0'}) = 2\pi (c^+_{p_0})^{-1}\delta(p_0 - p_0')$. An orthonormal basis is then given by

$$\begin{aligned}
\tilde{\xi}^+_{p_0} &:= \sqrt{c^+_{p_0}}\xi^+_{p_0} && (71)\\
\tilde{\xi}^-_{p_0} &:= \sqrt{c^-_{p_0}}\left(\xi^-_{p_0} - \frac{(\xi^-_{p_0}, \xi^+_{p_0})}{(\xi^+_{p_0}, \xi^+_{p_0})}\xi^+_{p_0}\right), && (72)
\end{aligned}$$

with $\sqrt{c^-_{p_0}}$ such that $(\tilde{\xi}^-_{p_0}, \tilde{\xi}^-_{p_0}) = 2\pi\delta(p_0 - p_0')$. From equation (55) we find the response of the detector as

$$P^{\Omega_0,p_0}(e \to g) = \frac{2\lambda^2}{m_0\Omega_0}\int_{-\infty}^{\infty} d\tau d\tau'\, e^{i(\Omega_0 - p_0)(\tau-\tau')}\left(|\tilde{\xi}^+_{p_0}(z)|^2 + |\tilde{\xi}^-_{p_0}(z)|^2\right). \quad (73)$$

If the complex structure $J_{\partial M}$ is unitary in the sense of our definition at the end of section 3.2.2 we have that if a function $e^+_{p_0} \in L^{\mathbb{C}}_{\tilde{M}}$ is an eigenfunction of J_{Σ_r} with $J_{\Sigma_r} e^+_{p_0} = -ie^+_{p_0}$ we get $J_{\Sigma_l} e^+_{p_0} = ie^+_{p_0}$, where the sign change with respect to the definition in 3.2.2 comes up because of the reverse orientation of Σ_l. We find that in this case $\tilde{\xi}^-_{p_0} = \overline{\sqrt{c^+_{p_0}}\xi^+_{-p_0}}$ and

$$c^+_{p_0} = ip_1\left(c_a(p_0)\overline{c_b(p_0)} - c_b(p_0)\overline{c_a(p_0)}\right) \quad (74)$$

and the response of the detector reduces to

$$\begin{aligned}
P^{\Omega_0,p_0}(e \to g) &= \frac{2\lambda^2}{m_0\Omega_0 p_1}\int_{-\infty}^{\infty} d\tau d\tau'\, e^{i(\Omega_0 - p_0)(\tau-\tau')}\\
&\qquad \sum_{p_0' = \pm p_0} \frac{|c_a(p_0')\cosh(p_1' z) + c_b(p_0')\sinh(p_1' z)|^2}{i\left(c_a(p_0)\overline{c_b(p_0)} - c_b(p_0)\overline{c_a(p_0)}\right)}.
\end{aligned}$$

If we require the complex structure to be invariant under time reversal which is a symmetry of our physical configuration then $c_a(-p_0) = c_a(p_0)$ and similarly for $c_b(p_0)$, and we find that

$$P^{\Omega_0,p_0}(e \to g) = \frac{\lambda^2}{m_0\Omega_0 p_1}\int_{-\infty}^{\infty} d\tau d\tau'\, e^{i(\Omega_0 - p_0)(\tau-\tau')}\frac{|c_a(p_0)\cosh(p_1 z) + c_b(p_0)\sinh(p_1 z)|^2}{\frac{i}{4}\left(c_a(p_0)\overline{c_b(p_0)} - c_b(p_0)\overline{c_a(p_0)}\right)},. \quad (75)$$

We want to have (75) proportional to the expression from the canonical model in (65) this determines $c_a(p_0)/c_b(p_0)$ and hence the complex structure: Since

$$\begin{aligned}
& |c_a(p_0)\cosh p_1 z + c_b(p_0)\sinh p_1 z|^2\\
= \ & \frac{1}{2}\left(|c_a(p_0)|^2 - |c_b(p_0)|^2 + (|c_a(p_0)|^2 + |c_b(p_0)|^2)\cosh 2p_1 z\right.\\
& \left. + 2|c_a(p_0)||c_b(p_0)|\cos(\phi_a - \phi_b)\sinh 2p_1 z\right) \quad (76)
\end{aligned}$$

with $c_a(p_0) = |c_a(p_0)|e^{i\phi_a(p_0)}$ and $c_b(p_0) = |c_b(p_0)|e^{i\phi_b(p_0)}$. Comparing with (65) this leads immediately to the condition $\phi_a(p_0) - \phi_b(p_0) \in \{\frac{\pi}{2}, \frac{3\pi}{2}\}$. This is determined to $\phi_a(p_0) - \phi_b(p_0) = \frac{\pi}{2}$ by the requirement of $J_{\partial M}$ being positive. We obtain two further conditions:

$$\begin{aligned}
|c_a(p_0)|^2 - |c_b(p_0)|^2 &\propto \frac{p_1}{p} - \frac{p}{p_1}\\
|c_a(p_0)|^2 + |c_b(p_0)|^2 &\propto \left(\frac{p_1}{p} + \frac{p}{p_1}\right)\cosh 2p_1\xi \quad (77)
\end{aligned}$$

which leads to

$$\frac{|c_a(p_0)|}{|c_b(p_0)|} = \left(\frac{\left(\frac{p_1}{p}-\frac{p}{p_1}\right)+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1\xi}{\left(\frac{p_1}{p}-\frac{p}{p_1}\right)-\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1\xi}\right)^{1/2} \tag{78}$$

We find that the complex structure leading to the same space dependence in (75) as in the canonical result (65) can be given by

$$\begin{aligned} c_a(p_0) &= \left(\left(\frac{p_1}{p}-\frac{p}{p_1}\right)+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1\xi\right)^{1/2} \\ c_b(p_0) &= i\left(\left(\frac{p_1}{p}-\frac{p}{p_1}\right)-\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1\xi\right)^{1/2} \end{aligned} \tag{79}$$

with $p=\sqrt{np_0^2-m^2}$ and N_{p_0} as given in equation (62). With that definition we obtain

$$P^{\Omega_0,\mathrm{out}}(e\to g)=\frac{\lambda^2}{m_0\Omega_0 p_1}|N_{\Omega_0}|^2\left(\left(\frac{p_1}{p}-\frac{p}{p_1}\right)+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1 z\,\cosh 2p_1\xi\right)\delta(0)^2\,, \tag{80}$$

which differs from the expression in equation (65) by a factor $\sqrt{n}\frac{p_1}{p}$ which comes up because of the definition of a particle as a normalized state and the difference of the normalization to the canonical case. We find that the condition to recover the expression in equation (65) we got from the canonical model together with the unitarity condition and the claim that the complex structure has all the symmetries of the classical problem determines the complex structure completely.

The necessity to require the unitarity of $J_{\partial M}$ can be seen by considering the complex structure given by eigenfunctions as in (70) with

$$\begin{aligned} c_{a;\Sigma_l}(p_0) &= \cosh(p_1\xi)+i\frac{p}{p_1}\sinh(p_1\xi) \\ c_{b;\Sigma_l}(p_0) &= \sinh(p_1\xi)+i\frac{p}{p_1}\cosh(p_1\xi)\,, \end{aligned}$$

and

$$\begin{aligned} c_{a;\Sigma_r}(p_0) &= \cosh(p_1\xi)-i\frac{p}{p_1}\sinh(p_1\xi) \\ c_{b;\Sigma_r}(p_0) &= -\sinh(p_1\xi)+i\frac{p}{p_1}\cosh(p_1\xi)\,. \end{aligned}$$

This gives a non-unitary complex structure which can be shown to recover the canonical result (65) from equation (73) as we prove in appendix A.4.

5.4.4. Spacelike asymptotics of the complex structure

In the foregoing section, we presented two different complex structures that lead to a split of the one particle Hilbert space into an orthogonal direct sum of linear subspaces that can be interpreted as outgoing states leading to a total probability $P^{\mathrm{out}}(g\to e)=0$ and ingoing states leading to $P^{\mathrm{in}}(g\to e)$ being proportional to the result in equation (65). The first of these two complex structures was unitary and the second not. In this section we investigate the asymtotics of these configurations when taking the boundaries Σ_l and Σ_r of the region M to spacelike infinity while evolving the complex structures on the hypersurfaces unitarily[10], i.e. having given the evolution map T from a timelike hyperplane Σ_I in the region I to Σ_l we give the corresponding complex structure as $J_{\Sigma_I} = T^{-1} J_{\Sigma_r} T$ etc.. This is easily performed by extending the eigenfunctions $e^+_{p_0;\Sigma_l}$ and $e^+_{p_0;\Sigma_r}$ of the complex structure to the regions I and III respectively. We obtain

---

[10] That was already done for the case of empty $3+1$-dimensional Minkowski space and a region bounded by a timelike hypercylinder in [CO08a, CO08b].

$$e^+_{p_0;\Sigma_r}(x_0,x_1) = \begin{cases} e_{p_0;\Sigma_r}(x_1) e^{-ip_0x_0} & \text{for} \quad -\xi \leq x_1 \leq \xi \\ \Big(e_{p_0;\Sigma_r}(\xi) \cos p(x_1-\xi) + \\ +\frac{1}{p} \left(\frac{d}{dx_1} e_{p_0;\Sigma_r}(x_1) \right)\Big|_{x_1=\xi} \sin p(x_1-\xi) \Big) e^{-ip_0x_0} & \text{for} \quad \xi < x_1 \,, \end{cases} \tag{81}$$

where $e_{p_0;\Sigma_r}(x_1) := c_{a;\Sigma_r}(p_0) \cosh(p_1 x_1) + c_{b;\Sigma_r}(p_0) \sinh(p_1 x_1)$ and

$$e^+_{p_0;\Sigma_l}(x_0,x_1) = \begin{cases} \Big(e_{p_0;\Sigma_l}(-\xi) \cos p(x_1+\xi) + \\ +\frac{1}{p} \left(\frac{d}{dx_1} e_{p_0;\Sigma_l}(x_1) \right)\Big|_{x_1=-\xi} \sin p(x_1+\xi) \Big) e^{-ip_0x_0} & \text{for} \quad x_1 < -\xi \\ e_{p_0;\Sigma_l}(x_1) e^{-ip_0x_0} & \text{for} \quad -\xi \leq x_1 \leq \xi \,, \end{cases} \tag{82}$$

where $e_{p_0;\Sigma_l}(x_1) := c_{a;\Sigma_l}(p_0) \cosh(p_1 x_1) + c_{b;\Sigma_l}(p_0) \sinh(p_1 x_1)$. We can compare this to the complex structure for propagating waves in equation (31) defined by the pressure condition in analogy to the AM-condition for spacelike hypersurfaces. We find that the unitary complex structure does not and the non-unitary complex structure does coincide with (31) when unitarily propagated to the regions I and III[11].

Note also that, as in the canonical theory, the extensions to the regions I and III of $e^+_{p_0;\Sigma_l}$ and $e^+_{p_0;\Sigma_r}$ respectively are only smooth, but not analytic at $x_1 = \pm\xi$ since the spacetime metric g has a discontinuity at $-\xi$ respectively ξ. Thus in particular an arbitrary small region in spacetime will no longer automatically contain the complete information on all spacetime without recurse to the equations of motion. Mathematically it is this fact that allows us to study the dependence of the complex structure on the physics outside of the spacetime region.

6. Conclusions

We have seen that Ashtekar-Magnon like conditions and symmetry considerations can indeed determine the remaining quantization ambiguity, in the holomorphic language the complex structure, for the propagating sector of quantum field theory on timelike hyperplanes. However, the evanescent modes also present in the theory require different treatment and new physical input.

We showed that the transition rates of a coupled detector are an appropriate tool to determine the complex structure in the evanescent sector. Interestingly, we found that even if the spacetime region under consideration is isometric to a region in Minkowski space the complex structure depends on the physics outside the spacetime region. In this sense we cannot describe the physics in a spacetime region purely locally without considering the embedding of the region into a spacetime.

In order to study this physical situation we had to re-express the condition that we have a transition from the vacuum to the one particle sector in GBF language. We saw that considering energy fluxes was a viable method in our case. However, we only studied these for highly symmetric and simple cases. An interesting open problem then is to understand the full quantized energy momentum tensor and it's detailed dependence on the complex structure. This would allow us to handle a much richer class of physical phenomena.

In the cases considered we were able to satisfy the physical conditions using a unitary complex structure, but it seems unlikely that this will be possible in general. The extension of the theory into the asymptotic regime also suggests that the unitary structure is not necessarily the physically preferred one, it is a non-unitary structure that matches up with the one chosen by the Ashtekar-Magnon like conditions. It is also unclear a priori that we can always model the physics in a spacetime region by only adjusting the complex structure.

---

[11] The non-unitary complex structure was chosen by the authors such that it fulfills this property in the first place. The interesting fact is that it leads to the correct detection rates.

Finally the result here should also be contrasted with the classic result of Reeh and Schlieder [Ree61, Sch65, Sum08] in algebraic quantum field theory. They showed that the vacuum state is a highly non-local structure. To compare to the language of algebraic quantum field theory we could describe the physical situation considered above as such: Consider two different, static, spacetimes, which coincide in a certain region. If we model the vacuum state of the second spacetime in the first one by applying operators localized outside of the region where the spacetimes coincide, we will generically change the vacuum state inside the coinciding region, too. The restrictions of the vacuum states to the coinciding spacetime region do not coincide, which is exactly what we found. As the response of the detector to evanescent modes is exponentially decaying away from the dielectric boundary, we have in some sense that, as in AQFT, the non-locality is exponentially small.

Acknowledgements

DR thanks Robert Oeckl, Daniele Colosi and Max Dohse for very helpful discussions, especially during a visit at the UNAM Campus Morelia. This research stay was funded by CONACYT grant 49093. The authors thank Robert Oeckl for reading the manuscript and making very helpful remarks. The work of RB and DR has been supported by the International Max Planck Research School for Geometric Analysis, Gravitation and String Theory.

A. Appendix

A.1. Proof of statement 1

In this section we show that for a timelike hyperplane Σ in Minkowski space

$$2\omega(\phi, n^\mu \partial_\mu \phi) = -2 \int_\Sigma d\sigma_\mu \, T^{\mu\nu} n_\nu \qquad \forall \phi \in L_\Sigma$$

is always satisfied. $T_{\mu\nu}$ is the energy-momentum tensor

$$T^{\mu\nu} = g^{\mu\alpha} g^{\mu\beta} \partial_\alpha \phi \partial_\beta \phi - \frac{1}{2} g^{\mu\nu} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - m^2 \phi^2)$$

and n is the gradient covector field to Σ with respect to the spacelike foliation. n is normalized to give $g(n, n) = -1$. We choose global coordinates $(x_0, x_1, \tilde{x})$ such that the metric takes everywhere the form $g = \text{diag}(1, -1, -1, -1)$. Without loss of generality we assume the timelike hyperplane Σ to be an equal x_1 hyperplane at $x_1 = \xi$ and to be oriented in positive x_1 direction. The measure $d\sigma_\mu$ is then given as $d\sigma_\mu = dx_{||} \, n_\mu$ with $dx_{||} := dx_0 dx_2 dx_3$. We write $n^\mu \partial_\mu = \partial_\perp$. Write out the LHS:

$$2\omega(\phi, n^\mu \partial_\mu \phi) = \int_\Sigma dx_{||} \, (\phi \partial_\perp^2 \phi - \partial_\perp \phi \partial_\perp \phi) \tag{83}$$

Let h^{ij} be the metric on Σ. Then we can write the equations of motion as $0 = g^{\mu\nu} \partial_\mu \partial_\nu \phi + m^2 \phi = (-\partial_\perp^2 + h^{ij} \partial_i \partial_j + m^2) \phi$, and plugging this into (83) and using integration by parts yields

$$\begin{aligned}
2\omega(\phi, n^\mu \partial_\mu \phi) &= \int_\Sigma dx_{||} \, \left(\phi (h^{ij} \partial_i \partial_j + m^2) \phi - \partial_\perp \phi \partial_\perp \phi \right) \\
&= -\int_\Sigma dx_{||} \, \left(h^{ij} \partial_i \phi \partial_j \phi - m^2 \phi^2 + \partial_\perp \phi \partial_\perp \phi \right) \\
&= -\int_\Sigma dx_{||} \, \left(g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - m^2 \phi^2 + 2 \partial_\perp \phi \partial_\perp \phi \right) \\
&= -2 \int_\Sigma dx_{||} \, T_{\perp\perp} = -2 \int_\Sigma d\sigma_\mu \, T^{\mu\nu} n_\nu .
\end{aligned}$$

This finishes the proof.

A.2. Proof of statement 2

We give the proof that in the case of Klein-Gordon theory on timelike hyperplanes Σ in $3+1$ Minkowski space, the requirement of J being compatible with ω is only satisfied by one complex structure out of the class of complex structures that may be written using a directional derivative with respect to some translation Killing vector field η of Minkowski space.

J is compatible with ω iff $g(\phi, \phi') = \omega(\phi, J\phi')$ is positive definite and $\omega(J\phi, J\phi') = \omega(\phi, \phi')$.

The symplectic structure in this case becomes

$$2\omega(\phi, \phi') = \int_\Sigma dx_{||}(\phi \partial_1 \phi' - \phi' \partial_1 \phi).$$

In this notation, we use global coordinates (x_0, x_1, x_2, x_3) such that the metric takes the form $g = \mathrm{diag}(1, -1, -1, -1)$ and $dx_{||} := dx_0 dx_2 dx_3$ is the measure for the integration in the hypersurface. Without loss of generality we consider the hyperplane $\Sigma_0 = \{x_1 = 0\}$. We shall parameterize the solution space of propagating waves L^p_Σ in the following manner:

$$\phi = \int \frac{dk_{||}}{(2\pi)^3} \frac{1}{2k_1} \left[\varphi(k) e^{ik\cdot x} + \overline{\varphi(k)} e^{-ik\cdot x}\right],$$

where $dk_{||} = dk_0 dk_2 dk_3$, $k\cdot x = -k_1 x_1 + k_{||}\cdot x_{||}$ with $k_{||}\cdot x_{||} = k_0 x_0 - k_2 x_2 - k_3 x_3$, $\varphi(k) \equiv \varphi(k_0, k_2, k_3)$ is a complex function on $\mathbb{R}^3$ and it is understood that $k_1 = \sqrt{|k_0^2 - k_2^2 - k_3^2 - m^2|}$. We shall assume (this assumption will prove to be correct later) that the plane waves are eigenfunctions of the complex structure J, that is

$$J e^{\pm ik\cdot x} = j_k^\pm \, i e^{\pm ik\cdot x}. \tag{84}$$

where $j_k^\pm \in \{-1, 1\}$ are yet to be determined (as functions of k_0, k_2 and k_3, possibly.). Now we can compute

$$\begin{aligned}
2\omega(\phi, J\phi) \;&=\; \int dx_{||}(\phi \partial_1 J\phi - J\phi \partial_1 \phi) \\
&=\; \int dx_{||} \int \frac{dk_{||}}{(2\pi)^3} \frac{1}{2k_1} \int \frac{dk'_{||}}{(2\pi)^3} \frac{1}{2k'_1} \Big(\Big[\varphi(k) e^{ik\cdot x} + \overline{\varphi(k)} e^{-ik\cdot x}\Big] \times \\
&\quad \Big[-i^2 k'_1 j^+_{k'} \varphi(k') e^{ik'\cdot x} + i^2 k'_1 j^-_{k'} \overline{\varphi(k')} e^{-ik'\cdot x}\Big] - \\
&\quad \Big[i\varphi(k) j_k^+ e^{ik\cdot x} + i\overline{\varphi(k)} j_k^- e^{-ik\cdot x}\Big] \times \\
&\quad \Big[-ik'_1 \varphi(k') e^{ik'\cdot x} + ik'_1 \overline{\varphi(k')} e^{-ik'\cdot x}\Big]\Big)
\end{aligned}$$

Now write $k_\mu x^\mu = k_1 x_1 + k_{||} \cdot x_{||}$, and use that

$$\int dx_{||} e^{-i(k_{||} \pm k'_{||}) x_{||}} = (2\pi)^3 \delta(k_{||} \pm k'_{||}),$$

which will also put $k'_1 = k_1$. Look now at all terms involving two $\varphi(k)$'s after integrating out $x_{||}$. We obtain

$$\int \frac{dk_{||}}{(2\pi)^3} \frac{1}{(2k_1)^2} k_1 \left[\varphi(k)\varphi(-k) j^+_{-k} e^{2ik_1 x_1} - \varphi(k)\varphi(-k) j^+_k e^{2ik_1 x_1}\right] = 0$$

which gives zero after substituting $k_{||} \to -k_{||}$ in the second term. The same happens with all contributions involving two $\overline{\varphi(k)}$'s. One is then left with

$$2\omega(\phi, J\phi) = \int \frac{dk_{||}}{(2\pi)^3} \frac{1}{2k_1} \frac{1}{2} \left(\varphi(k)\overline{\varphi(k)} + \overline{\varphi(k)}\varphi(k) \right) \left[j_k^+ - j_k^- \right].$$

The condition that $g(\phi, \phi) = \omega(\phi, J\phi)$ is positive definite then indicates that $j_k^+ - j_k^-$ should never be negative. Let us now focus on a certain class of complex structures $J_\eta = \frac{\partial_\eta}{\sqrt{-\partial_\eta^2}}$ parameterized by a vector η. Then it's clear that

$$\partial_\eta e^{\pm ik\cdot x} = \pm i\eta \cdot k\, e^{\pm ik\cdot x}.$$

In particular, this means that the plane waves are eigenfunctions of J, as we assumed earlier. And now keeping in mind how the j's are defined in (84), we obtain

$$j_k^+ = \operatorname{sgn}(\eta \cdot k) \tag{85}$$

$$j_k^- = -\operatorname{sgn}(\eta \cdot k) = -j_k^+ \tag{86}$$

This gives

$$\omega(\phi, J\phi) = \int \frac{dk_{||}}{(2\pi)^3} \frac{1}{2k_1}\, \varphi(k)\overline{\varphi(k)}\, \operatorname{sgn}(\eta \cdot k).$$

This is positive definite iff

$$\eta \cdot k = \eta_0 k_0 - \eta_1 \sqrt{|k_0^2 - k_2^2 - k_3^2 - m^2|} - \eta_2 k_2 - \eta_3 k_3 \geq 0$$

holds for all k_0, k_2, k_3 such that $k_0^2 - k_2^2 - k_3^2 - m^2 \geq 0$. In particular, it has to hold for those k_0, k_2, k_3 for which $k_0^2 - k_2^2 - k_3^2 - m^2 = 0$ [12]. Then the positive definiteness condition reduces to

$$\eta_0 k_0 - \eta_2 k_2 - \eta_3 k_3 \geq 0. \tag{87}$$

Now of course $k_0 = \pm\sqrt{k_2^2 + k_3^2 + m^2}$ depends on k_2 and k_3, but we still have a sign freedom in k_0, k_2 and k_3. This means that for any given (η_0, η_2, η_3), we can find (k_0, k_2, k_3) such that the LHS of (87) becomes strictly negative, with the exception being $\eta_0 = \eta_2 = \eta_3 = 0$. This leaves η_1 to be negative and determines η as the direction perpendicular to the hypersurface, and thus J. One can check that this J indeed satisfies $\omega(\phi, \phi') = \omega(J\phi, J\phi')$.

A.3. Orthonormality of the functions in equation (61)

First of all, we show that solutions of different energy are orthogonal in any case. This we see from the time translation invariance of the symplectic structure:

$$0 = \partial_0 \omega_{x_0}(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r) = i(p_0 - p_0')\omega_{x_0}(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r) = (p_0 - p_0')(\phi_{p_0}^s, \phi_{p_0'}^r)$$

which means for $p_0 \neq p_0'$ that $(\phi_{p_0}^s, \phi_{p_0'}^r)$ must vanish. Now, equation (63) can be derived using the continuity equation $\partial_\mu j^\mu(\phi, \phi') = 0$. We define $j^\mu(\phi, \phi') := \sqrt{g}\, g^{\mu\nu}(\phi\partial_\nu\phi' - \phi'\partial_\nu\phi)$, and then

$$j^0(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r) = i\frac{1}{p_0 - p_0'}\partial_1 j^1(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r) \tag{88}$$

which leads to

$$\begin{aligned} \omega_{x_0}(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r) &= \frac{i}{2}\lim_{\sigma\to\infty} \frac{1}{p_0 - p_0'} j^1(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r)(x)\Big|_{x_1=-\sigma}^{x_1=\sigma} \\ &= \frac{i}{2}\lim_{\sigma\to\infty} \frac{n}{p - p'}\frac{p_0 + p_0'}{p + p'} j^1(\overline{\phi_{p_0}^s}, \phi_{p_0'}^r)(x)\Big|_{x_1=-\sigma}^{x_1=\sigma}. \end{aligned}$$

---

[12] That we can make this argument is of course due to the hyperplane Σ being timelike. For spacelike hyperplanes, the positive definiteness condition for g only bounds the vector η to the light cone.

Then, we plug in the functions given in (61) and neglect all terms that come with the factor $p+p'$ since they vanish in the limit $\sigma \to 0$. Using the identities $\overline{C_{p_0}}N_{p_0}+C_{p_0}\overline{N_{p_0}}=0$ and $|C_{p_0}|^2+|N_{p_0}|^2=1$ and the representation of the delta function

$$\lim_{\sigma\to\infty}\frac{\sin(p-p')\sigma}{p-p'}=\pi\delta(p-p')$$

we arrive at

$$\begin{aligned}\omega_{x_0}(\overline{\phi^+_{p_0}},\phi^-_{p'_0}) &= 0\\ \omega_{x_0}(\overline{\phi^+_{p_0}},\phi^+_{p'_0}) &= i\sqrt{n}2\pi p_0\delta(p-p')\end{aligned}$$

which leads to equation (63).

A.4. Recovering the space dependence of the canonical result with a non-unitary complex structure

First, we find that we can express the eigenfunctions of the complex structures of J_{Σ_l} and J_{Σ_r} in terms of each other by the relation

$$\left(\frac{e^+_{p_0,\Sigma_l}}{e^+_{-p_0,\Sigma_l}}\right)=\begin{pmatrix} A(p_0) & B(p_0)\\ \overline{B(p_0)} & \overline{A(p_0)}\end{pmatrix}\left(\frac{e^+_{p_0,\Sigma_r}}{e^+_{-p_0,\Sigma_r}}\right)=:T\left(\frac{e^+_{p_0,\Sigma_r}}{e^+_{-p_0,\Sigma_r}}\right), \tag{89}$$

where

$$\begin{aligned}A(p_0) &= -\frac{i}{2}\left(\frac{p_1}{p}-\frac{p}{p_1}\right)\sinh 2p_1\xi+\cosh 2p_1\xi\\ B(p_0) &= \frac{i}{2}\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\sinh 2p_1\xi\end{aligned}$$

and we find that

$$T^{-1}=\begin{pmatrix}\overline{A(p_0)} & -B(p_0)\\ -\overline{B(p_0)} & A(p_0)\end{pmatrix}. \tag{90}$$

From that we calculate for $\xi^+_{p_0}=e^+_{p_0,\Sigma_r}$ and $\xi^-_{p_0}=\overline{e^+_{-p_0,\Sigma_r}}$

$$\begin{aligned}(\xi^+_{p_0},\xi^+_{p'_0})&=g_{\Sigma_l}(\xi^+_{p_0},\overline{\xi^+_{p'_0}})+g_{\Sigma_r}(\xi^+_{p_0},\overline{\xi^+_{p'_0}})\\ =(|A(p'_0)|^2+|B(p'_0)|^2+1)g_{\Sigma_r}(\xi^+_{p_0},\overline{\xi^+_{p'_0}})&=(|A(p_0)|^2+|B(p_0)|^2+1)2\pi p\delta(p_0-p'_0)\end{aligned}$$

and the same way

$$\begin{aligned}(\xi^-_{p_0},\xi^+_{p'_0}) &= g_{\Sigma_l}(\xi^-_{p_0},\overline{\xi^+_{p'_0}})+g_{\Sigma_r}(\xi^-_{p_0},\overline{\xi^+_{p'_0}})\\ &= -2A(p_0)\overline{B(p_0)}\,g_{\Sigma_r}\left(\overline{e^+_{-p_0,\Sigma_r}},e^+_{-p_0,\Sigma_r}\right)=-2A(p_0)\overline{B(p_0)}2\pi p\delta(p_0-p'_0)\end{aligned}$$

which leads us to

$$\begin{aligned}(c^+_{p_0})^{-1} &= \frac{p}{p_1}(|A(p_0)|^2+|B(p_0)|^2+1)\\ &= \frac{p}{4p_1}\left(6-\frac{p_1^2}{p^2}-\frac{p^2}{p_1^2}+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)^2\cosh 4p_1\xi\right).\end{aligned}$$

Furthermore, we define

$$g_{p_0} := -2\frac{p}{p_1}A(p_0)\overline{B(p_0)} = \frac{p}{2p_1}\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\sinh 2p_1\xi\left(\left(\frac{p_1}{p}-\frac{p}{p_1}\right)\sinh 2p_1\xi + 2i\cosh 2p_1\xi\right)$$

and obtain

$$(c^-_{p_0})^{-1} = (c^+_{p_0})^{-1}\left(1-(c^+_{p_0})^2|g_{p_0}|^2\right) = 2\frac{p}{p_1}\,. \tag{91}$$

With the definition in equation (71) we have

$$\begin{aligned}
\tilde{\xi}^+_{p_0} &:= \sqrt{c^+_{p_0}}\xi^+_{p_0} \\
\tilde{\xi}^-_{p_0} &:= \sqrt{c^-_{p_0}}\left(\xi^-_{p_0} - c^+_{p_0}g_{p_0}\xi^+_{p_0}\right) \\
&= \sqrt{\frac{p_1}{2p}}\left(\cosh 2p_1\xi + \frac{i}{2}\left(\frac{p_1^2}{p^2}-\frac{p^2}{p_1^2}\right)\sinh 2p_1\xi\right)^{-1}\times \\
&\quad\times\left(\cosh p_1(x+\xi) - i\frac{p}{p_1}\sinh p_1(x+\xi)\right)
\end{aligned}$$

and arrive at

$$|\tilde{\xi}^+_{p_0}|^2 + |\tilde{\xi}^-_{p_0}|^2 = \frac{1}{2}|N_{p_0}|^2\left(\left(\frac{p_1}{p}-\frac{p}{p_1}\right)+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)\cosh 2p_1x_1\,\cosh 2p_1\xi\right)$$

with

$$|N_{p_0}|^{-2} = \frac{1}{8}\left(6-\frac{p_1^2}{p^2}-\frac{p^2}{p_1^2}+\left(\frac{p_1}{p}+\frac{p}{p_1}\right)^2\cosh 4p_1\xi\right) \tag{92}$$

which shows that we recover the canonical result.

References

[AM75] A. Ashtekar and A. Magnon, *Quantum fields in curved space-times*, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences **346**(1646), 375 (1975).

[Ati89] M. Atiyah, *Topological quantum field theories*, Inst. Hautes Etudes Sci. Publ. Math. **68**, 175–186 (1989).

[CDO11] D. Colosi, M. Dohse and R. Oeckl, *S-Matrix for AdS from General Boundary QFT*, (2011), 1112.2225.

[CM71] C. Carniglia and L. Mandel, *Quantization of evanescent electromagnetic waves*, Physical Review D **3**(2), 280 (1971).

[CMD72] C. Carniglia, L. Mandel and K. Drexhage, *Absorption and emission of evanescent photons*, JOSA **62**(4), 479–486 (1972).

[CO08a] D. Colosi and R. Oeckl, *S-matrix at spatial infinity*, Phys. Lett. **B665**, 310–313 (2008), 0710.5203.

[CO08b] D. Colosi and R. Oeckl, *Spatially asymptotic S-matrix from general boundary formulation*, Phys. Rev. **D78**, 025020 (2008), 0802.2274.

[CO09] D. Colosi and R. Oeckl, *States and amplitudes for finite regions in a two- dimensional Euclidean quantum field theory*, J. Geom. Phys. **59**, 764–780 (2009), 0811.4166.

[CO11] D. Colosi and R. Oeckl, *On Unitary Evolution in Quantum Field Theory in Curved Spacetime*, Open Nucl. Part. Phys. J. **4**, 13–20 (2011), 0912.0556.

[EMRV99] A. Echeverria-Enriquez, M. C. Munoz-Lecanda, N. Roman-Roy and C. Victoria-Monge, *Mathematical Foundations of Geometric Quantization*, ArXiv Mathematical Physics e-prints (April 1999), arXiv:math-ph/9904008.

[FD99] J. Fischer and T. Dray, *A New Look at the Ashtekar–Magnon Energy Condition*, General Relativity and Gravitation **31**(4), 511–526 (1999).

[LH07] S. Lin and B. Hu, *New Insights into Uniformly Accelerated Detector in a Quantum Field*, Foundations of Physics **37**(4), 480–490 (2007).

[Oec05] R. Oeckl, *States on timelike hypersurfaces in quantum field theory*, Phys. Lett. **B622**, 172–177 (2005), hep-th/0505267.

[Oec06] R. Oeckl, *General boundary quantum field theory: Timelike hypersurfaces in Klein-Gordon theory*, Phys. Rev. **D73**, 065017 (2006), hep-th/0509123.

[Oec07] R. Oeckl, *Probabilites in the general boundary formulation*, J. Phys. Conf. Ser. **67**, 012049 (2007), hep-th/0612076.

[Oec08] R. Oeckl, *General boundary quantum field theory: Foundations and probability interpretation*, Adv. Theor. Math. Phys. **12**, 319–352 (2008), hep-th/0509122.

[Oec10] R. Oeckl, *Holomorphic quantization of linear field theory in the general boundary formulation*, (2010), 1009.5615.

[Oec11] R. Oeckl, *The Schrödinger representation and its relation to the holomorphic representation in linear and affine field theory*, (2011), 1109.5215.

[Oec12a] R. Oeckl, *Affine holomorphic quantization*, J.Geom.Phys. **62**, 1373–1396 (2012), 1104.5527.

[Oec12b] R. Oeckl, Observables in the General Boundary Formulation, in *Quantum Field Theory and Gravity*, edited by F. Finster, O. Müller, M. Nardmann, J. Tolksdorf and E. Zeidler, pages 137–156, Springer Basel, 2012.

[Oec12c] R. Oeckl, *Schrödinger-Feynman quantization and composition of observables in general boundary quantum field theory*, (2012), 1201.1877.

[Ree61] Reeh H., Schlieder S., Bemerkungen zur Unitäräquivalenz von Lorentzinvarianten Feldern, 1961, Nuovo Cimento **22**, 1051.

[Sch65] S. Schlieder, *Some remarks about the localization of states in a quantum field theory.*, Commun. Math. Phys. **1**, 265–280 (1965).

[Sum08] S. J. Summers, *Yet More Ado About Nothing: The Remarkable Relativistic Vacuum State*, ArXiv e-prints (February 2008), 0802.1854.

[Woo97] N. Woodhouse, *Geometric Quantization*, Oxford Mathematical Monographs, Clarendon Press, 1997.